

FLAT CENTRAL DENSITY PROFILES FROM SCALAR FIELD DARK MATTER HALOS

Argelia Bernal,^{1,3} Tonatiuh Matos,^{1,3} and Darío Núñez^{2,3}

Received 2006 May 24; accepted 2008 January 10

RESUMEN

El campo escalar con un potencial cosh se comporta en el régimen lineal justo como la materia oscura fría (CDM). Así, la hipótesis de materia oscura escalar (SFDM) predice la misma formación de estructura que CDM. Los parámetros libres del modelo SFDM se fijan con observaciones cosmológicas. En un trabajo anterior mostramos que con dichos valores, la SFDM se colapsa y forma objetos estables con masa del orden de $10^{12} M_{\odot}$. En este trabajo utilizamos soluciones analíticas para las ecuaciones de Einstein-Klein-Gordon en el caso plano y en el límite de campo débil y mostramos que el perfil de densidad de SFDM corresponde al de una densidad central casi plana y que coincide con el del modelo de CDM en una amplia zona de la región exterior. Este resultado podría resolver el problema de los picos de densidad de CDM sin ninguna hipótesis adicional, lo que apoya la viabilidad del modelo de SFDM.

ABSTRACT

Scalar fields endowed with a cosh potential behave in the linear regime, exactly as the cold dark matter (CDM) model. Thus, the scalar field dark matter (SFDM) hypothesis predicts the same structure formation as the CDM model. This means that CDM and SFDM are equivalent from the cosmological point of view. The free parameters of the SFDM model are determined by cosmological observations. In previous work we showed that if we use such parameters, the scalar field collapses forming stable objects with a mass around $10^{12} M_{\odot}$. In the present work we use analytical solutions of the flat and weak field limit of the Einstein-Klein-Gordon equations and show that the SFDM density profile corresponds to a halo with an almost flat central density and that it coincides with the CDM model in a broad outer region. This result could solve the problem of the density cusp DM halo in galaxies without any additional hypothesis, thus supporting the viability of the SFDM model.

Key Words: dark matter — galaxies: formation — galaxies: halos

1. INTRODUCTION

The Lambda Cold Dark Matter (Λ CDM) model has recently shown an enormous predictive power. It can explain the structure formation of the Universe, its accelerated expansion, the micro Kelvin fluctuation of the cosmic microwave background radiation, etc. Nevertheless, around this model some issues related to the formation of galaxies have arisen since

the time it was originally proposed and remain to date. The CDM paradigm predicts a density profile which corresponds to the Navarro-Frenk-White (NFW) profile (Navarro, Frenk, & White 1997) given by

$$\rho_{NFW} = \frac{\rho_0}{\frac{r}{r_0}(\frac{r}{r_0} + 1)^2}. \quad (1)$$

However, this profile seems to differ somewhat from the observed profiles of LSB galaxies. In this work we show that a flat central profile naturally arises within the scalar field dark matter hypothesis, implying that the central region of galaxies can distinguish between CDM and SFDM.

¹Departamento de Física, Centro de Investigación y de Estudios Avanzados del IPN, Mexico.

²Instituto de Ciencias Nucleares, Universidad Nacional Autónoma de México, Mexico.

³Part of the Instituto Avanzado de Cosmología (IAC), Mexico.

We work within the specific context of the so-called ‘strong, self-interacting scalar field dark matter’ (SFDM) hypothesis that has been developed by several authors (Guzmán & Matos 2000; Matos & Ureña-López 2000, 2001; Ureña-López, Matos, & Becerril 2002; Matos & Guzmán 2001; Alcubierre et al. 2002, 2003; Ureña-López 2002; Böhmer & Harko 2007; see also Peebles 2000). A first proposal of the SFDM hypothesis appeared in a couple of papers by Ji & Sin 1994. They took a massive scalar field and were able to fit observations from some galaxies, taken into account also the contribution of baryons. A key point in this work was the use of the so-called excited configurations in which the radial profile of the scalar fields has nodes. From that, they determined that the mass of the scalar field should be of order of $\sim 10^{-24}$ eV.

The next proposal appeared in a paper by Schunck (1998). He shows that a massless complex scalar field can be used as a dark matter model in galaxies to fit the rotation curves. In this model the internal frequency of the field plays the role of an adjustable parameter, and the radial profile of the scalar field also has nodes.

However, as pointed out in Guzmán & Ureña-López (2005; see also Guzmán & Ureña-López 2004), this last proposal cannot be realistic because a massless scalar field (whether real or complex as in Schunck 1998) cannot form a gravitationally bound configuration (see Seidel & Suen 1994).

The key idea of the SFDM scenario is that the dark matter responsible for structure formation in the Universe is a real scalar field, Φ , minimally coupled to Einstein gravity with the self-interaction parametrized by a potential energy of the form (see also Sahni & Wang 2000)

$$V(\Phi) = V_0 [\cosh(\lambda\sqrt{\kappa_0}\Phi) - 1], \quad (2)$$

where V_0 and λ are the only two free parameters of the model, $\kappa_0 = 8\pi G$, and we employ natural units $\hbar = c = 1$. The effective mass of the scalar field is given by $m_\Phi^2 = \kappa_0 V_0 \lambda^2$.

The advantage of the SFDM model is that it is insensitive to initial conditions and the scalar field behaves as CDM once it begins to oscillate around the minimum of its potential. In this case, it can be shown (see Matos & Ureña-López 2000, 2001) that the SFDM model is able to reproduce all the successes of the standard Λ CDM model above galactic scales.

Furthermore, it predicts a sharp cut-off in the mass power spectrum due to its quadratic nature,

thus explaining the observed dearth of dwarf galaxies, in contrast with the possible excess predicted by high resolution N-body simulations with standard CDM (see Matos & Ureña-López 2001).

The best-fit model to the cosmological data can be deduced from the current densities of dark matter and radiation in the Universe and from the cut-off in the mass power spectrum that constrains the number of dwarf galaxies in clusters. The favored values for the two free parameters of the scalar field potential are found to be (Matos & Ureña-López 2001):

$$\begin{aligned} \lambda &\simeq 20, \\ V_0 &\simeq (3 \times 10^{-27} m_{Pl})^4, \end{aligned} \quad (3)$$

where $m_{Pl} \equiv G^{-1/2} \approx 10^{-5}g$ is the Planck mass.

This implies that the effective mass of the scalar field should be $m_\Phi \simeq 9.1 \times 10^{-52} m_{Pl} = 1.1 \times 10^{-23}$ eV.

Let us explain why we suspect that the scalar field could be the dark matter at galactic scales as well. There are three main reasons.

The first reason is that numerical simulations suggest that the critical mass for the case considered here, using the scalar potential (equation 2), and the parameters given by equation (3), is approximately (Alcubierre et al. 2002)

$$M_{crit} \simeq 0.1 \frac{m_{Pl}^2}{\sqrt{\kappa_0 V_0}} = 2.5 \times 10^{13} M_\odot. \quad (4)$$

This was a surprising result. The critical mass of the model shown in Matos & Ureña-López (2000, 2001) is of the same order of magnitude of the dark matter content of a standard galactic halo. Observe that the parameters of the model, equation (3), were fixed using cosmological observations. The surprising result consisted in the fact that using the same scalar field for explaining the dark matter at cosmological scales, it will always collapse with a preferred mass which corresponds to the halo of a real galaxy. Thus, this result is a prediction of the cosmological SFDM model for galaxy formation.

The second reason is that during the linear regime of cosmological fluctuations, the scalar field and a dust fluid, like CDM, behave in the same way. The density contrast in CDM and in the SFDM models evolve in exactly the same form, and thus both models predict the same large scale structure formation in the Universe (see Matos & Ureña-López 2001). The differences between the CDM and SFDM models begin to appear in the non-linear regime of structure formation, so that there will be differences in their predictions of galaxy formation.

The third reason is the topic of this work. A scalar field object (e.g. an oscillaton) contains a flat central density profile, as seems to be the case in galaxies.

In the case of the SFDM, the strong self-interaction of the scalar field results in the formation of solitonic objects called ‘oscillatons’, which have a mass of the order of a galaxy (see for example Ureña-López 2002; Ureña-López et al. 2002; Alcubierre et al. 2003; also Seidel & Suen 1991, 1994; Hawley & Choptuik 2000; Honda & Choptuik 2001). In this work we will show that these models contain an almost flat central density profile, *i.e.*, they do not exhibit the cusp density profiles characteristic of the standard CDM hypothesis.

Before starting with the description, we want to emphasize the fact that the scalar field has no interaction with the rest of the matter; thus, it does not follow the standard lines of reasoning for the particle-like candidates for dark matter. The scalar field was not thermalized, that is, the scalar field forms a Bose condensate, and thus behaves strictly as cold dark matter from the beginning.

The rest of the paper is organized as follows. In the next section we use the fact that Galaxies have a weak gravitational field and thus their space-time is almost flat. The main goal of this section is to study the physics provoking the flat behavior of the density profiles at the center of the oscillatons. Some results of this section intersect with those presented by Ji & Sin (1994), where they studied the behavior of the weak field limit of a complex scalar field. We remark that we do our analysis for a real scalar field and some differences do arise due to a different current conservation. In § 3 we present our contribution making an analysis of the Einstein-Klein-Gordon (EKG) equations in the relativistic weak field limit, and we solve them for the perturbed metric coefficients. Then, we compare these solutions with the ones obtained by solving numerically the complete EKG system and, using the whole potential (equation 2), we show that the relativistic weak field limit is indeed a very good approximation. In § 4 we compute the energy density of the scalar field obtained in § 3, and compare it with actual observations of LSB galaxies from which the density is inferred from the rotation curves, showing a good match in the external regions and a match at least similar in some of the internal regions, in any case, better than a fit with a cusp-like behavior of the density. Finally in § 5 we give our conclusions.

2. PHYSICS OF THE SCALAR FIELD. FLAT SPACE-TIME CASE

In this section we derive the physics of the scalar field in an analogous way as it was done for the complex scalar field by Ji & Sin (1994) (see also Lee & Koh 1996). In a normal dust collapse, as for example in CDM, there is in principle nothing to prevent that the dust matter collapses all the time. There is only a radial gravitational force that provokes the collapse, and to stop it one needs to invoke some virialization phenomenon. In the scalar field paradigm the collapse is different. The energy momentum tensor of the scalar field is

$$T_{\mu\nu} = \Phi_{,\mu}\Phi_{,\nu} - \frac{g_{\mu\nu}}{2} [\Phi^{,\alpha}\Phi_{,\alpha} + 2V(\Phi)] . \quad (5)$$

We will consider spherical symmetry, and work with the line element

$$ds^2 = -e^{2\nu} dt^2 + e^{2\mu} dr^2 + r^2 d\Omega^2 , \quad (6)$$

with $\mu = \mu(r, t)$ and $\nu = \nu(r, t)$, being this last function the Newtonian potential. The energy momentum tensor of the scalar field has then the components

$$-T_0^0 = \rho_\Phi = \frac{e^{-2\nu}\dot{\Phi}^2 + e^{-2\mu}\Phi'^2}{2} + V(\Phi) , \quad (7)$$

$$T_{01} = \mathcal{P}_\Phi = \dot{\Phi}\Phi' , \quad (8)$$

$$T_1^1 = p_r = \frac{e^{-2\nu}\dot{\Phi}^2 + e^{-2\mu}\Phi'^2}{2} - V(\Phi) , \quad (9)$$

$$T_2^2 = p_\perp = \frac{e^{-2\nu}\dot{\Phi}^2 - e^{-2\mu}\Phi'^2}{2} - V(\Phi) , \quad (10)$$

and also $T_3^3 = T_2^2$. These different components are identified as the energy density ρ_Φ , the momentum density \mathcal{P}_Φ , the radial pressure p_r and the angular pressure p_\perp . The integrated mass is defined by

$$M(x) = 4\pi \int_0^x \rho_\Phi(X) X^2 dX . \quad (11)$$

The radial and angular pressures are two natural components of the scalar field which stop the collapse, avoiding the cusp density profiles in the centers of the collapsed objects. This is the main difference between the normal dust collapse and the SFDM one. The pressures play an important role in the SFDM equilibrium. In order to see this, and considering that galaxies are almost flat, we conclude that the Newtonian approximation should be sufficient to describe the processes. In this section we will take the flat space-time approximation.

Thus, we study a massive oscillaton without self-interaction (*i.e.* with potential $V = 1/2(m_\Phi^2 \Phi^2)$),

in the Minkowski background ($\mu \sim \nu \sim 0$). Even though it is not a solution to the Einstein equations as we are neglecting the gravitational force provoked by the scalar field, the solution is analytic and it helps us to understand some features that appear in the non-flat oscillatons.

In a spherically symmetric space-time, the Klein-Gordon equation $\eta^{\alpha\beta}\partial_\alpha\partial_\beta\Phi - dV/d\Phi = 0$, (where $\eta^{\alpha\beta}\partial_\alpha\partial_\beta$ stands for the D'Alembertian), reads

$$\Phi'' + \frac{2}{r}\Phi' - m_\Phi^2\Phi = \ddot{\Phi}, \quad (12)$$

where over-dot denotes $\partial/\partial t$ and prime denotes $\partial/\partial r$. The exact general solution for the scalar field Φ is

$$\Phi(t, r) = \frac{e^{\pm ikr}}{r} e^{\pm i\omega t}, \quad (13)$$

and we obtain the dispersion relation $k^2 = \omega^2 - m_\Phi^2$. For $\omega > m_\Phi$ the solution is non-singular and vanishes at infinity. We will restrict ourselves to this case. It is more convenient to use trigonometric functions and to write the particular solution in the form

$$\Phi(t, x) = \Phi_0 \frac{\sin(x)}{x} \cos(\omega t), \quad (14)$$

where $x = kr$. It oscillates in harmonic manner in time. The scalar field can be considered to be confined to a finite region (see Ureña-López 2002; Ureña-López et al. 2002).

The analytic expression for the scalar field energy density derived from equation (14) is

$$\rho_\Phi = \frac{\Phi_0^2 k^2}{2x^2} \left[\left(\left(\frac{\sin(x)}{x} - \cos(x) \right)^2 - k^2 \sin^2(x) \right) \cos^2(\omega t) + \omega^2 \sin^2(x) \right], \quad (15)$$

which oscillates with a frequency $2\omega t$. Observe that close to the central regions of the object, the density of the oscillaton behaves like

$$\rho_\Phi \sim \frac{1}{2}\Phi_0^2 k^2 [\omega^2 - k^2 \cos^2(\omega t)] + O(x^2), \quad (16)$$

which implies that when $x \rightarrow 0$ the central density oscillates around a fixed value.

On the other hand, the asymptotic behavior when $x \rightarrow \infty$, is such that $\rho_\Phi \sim 1/x^2$, *i.e.*, far away from the center, in this approximation, the flat oscillaton density profile behaves like the isothermal one. The mass function oscillates around $M \sim x$, as usual for the galactic halos.




Fig. 1. The energy density profile for the Scalar Field Dark Matter model. The parameters used in this plot are $m_\Phi = 20$, $\omega = 20.001$, $\Phi_0^2 = 5 \times 10^{-2}$. The density is given in arbitrary units and r is given in kpc.

In order to understand what is happening within the object, observe that the KG equation can be rewritten in a more convenient form in terms of the energy density, as

$$\frac{\partial \rho_\Phi}{\partial t} - \frac{1}{r^2} \frac{\partial}{\partial r} (r^2 \mathcal{P}_\Phi) = 0. \quad (17)$$

This last equation has a clear interpretation: Since its form looks like the conservation equation, $\dot{\rho} + \nabla \cdot \vec{J} = 0$, equation (17) represents the conservation of the scalar field energy. It also tells us that there is a scalar field current given by

$$\begin{aligned} \vec{J}_\Phi &= -\mathcal{P}_\Phi \vec{r} \\ &= \Phi_0^2 \frac{k\omega}{2} \left[x \cos(x) - \sin(x) \right] \frac{\sin(x) \sin(2\omega t)}{x^3} \vec{r}. \end{aligned}$$

Observe that the quantity involved in this current is the scalar field momentum density (equation 8). Although the flux of scalar radiation at large distances does not vanish, there is no net flux of energy, as it can be seen by averaging the scalar current over a period of a scalar oscillation. We also see that the only transformation process is that of the scalar field energy density into the momentum density, and viceversa. For the realistic values (equation 3) this transfer is very small.

In Figure 1 we show the behavior of the SFDM density profile for a typical galaxy and in Figure 2 we show the comparison between the NFW, the isothermal and the SFDM density profiles for the same galaxy. Observe that the SFDM and NFW profiles remain very similar up to 10 kpc, then the SFDM profile starts to follow the isothermal one.




Fig. 2. Comparison between the energy density profile for the Scalar Field Dark Matter model with the NFW and the Isothermal models. The parameters for the isothermal model are $\rho_{\text{Iso}} = 0.3/(r^2 + 8^2)$ and for the NFW profile are $\rho_{\text{NFW}} = 10/(r(r+8)^2)$. The parameters used for the SFDM model are the same as in the previous figure.

In the next section we will see that if the gravitational force is taken into account, the oscillaton is more confined (see also Ureña-López 2002; Ureña-López et al. 2002; Alcubierre et al. 2003). The parameters used in the figures correspond to a medium size galaxy.

3. WEAK FIELD LIMIT EQUATIONS

In this section we derive a novel method for integrating the perturbed EKG equations and show that the solutions are in very good agreement with the numerical ones. This allows us to use these solutions to fit the rotation curves of several observed LSB galaxies.

Within general relativity, the evolution of the scalar and gravitational fields is governed by the coupled EKG equations, the last one appearing from the conservation of the energy-momentum tensor

$$R_{\alpha\beta} = \kappa_0 S_{\alpha\beta}, \quad (18)$$

$$T^{\alpha\beta}{}_{;\beta} = \Phi^{;\alpha}(\nabla^2 - m^2)\Phi = 0, \quad (19)$$

here $R_{\alpha\beta}$ is the Ricci tensor, $S_{\alpha\beta} = T_{\alpha\beta} - (1/2)g_{\alpha\beta}T^\lambda{}_\lambda$, and $\nabla^2 = (1/\sqrt{-g})\partial_\mu[\sqrt{-g}g^{\mu\nu}\partial_\nu]$ is the covariant D'Alembertian operator.

For simplicity, we continue to consider the non-static spherically symmetric case, given by equation (6). As usual in the weak field limit, we suppose the metric to be close to the Minkowski metric $\eta_{\alpha\beta}$

$$g_{\alpha\beta} = \eta_{\alpha\beta} + h_{\alpha\beta}, \quad (20)$$

where $|h_{\alpha\beta}| \ll 1$, then we will consider an expansion of the functions in the metric of the form

$$\begin{aligned} e^{2\nu(t,r)} &= 1 + \epsilon^2 2V(t,r) + O(\epsilon^4), \\ e^{2\mu(t,r)} &= 1 + \epsilon^2 2U(t,r) + O(\epsilon^4), \end{aligned} \quad (21)$$

where ϵ is an expansion parameter. We also consider that the spatial and time derivatives of the geometric quantities are regarded like

$$\frac{\partial}{\partial t} \sim \frac{\partial}{\partial r}. \quad (22)$$

Then to first order in ϵ^2 the Ricci tensor components are, respectively,

$$\begin{aligned} R_{tt} &= V_{,rr} - U_{,tt} + \frac{2}{r}V_{,r}, \\ R_{rr} &= U_{,tt} - V_{,rr} + \frac{2}{r}U_{,r}, \\ R_{tr} &= \frac{2}{r}U_{,t}. \end{aligned} \quad (23)$$

On the other hand, the source is computed in the flat space in this case as well, thus the scalar field satisfies to ϵ^2 order equation (12), as in the previous section.

It is important to emphasize that the relation (equation 22) is the lowest one in the geometric fields, but it does not consider small velocities for the sources. This is different from the Newtonian limit where the derivative relation for the scalar field is $\partial_r \sim \epsilon \partial_t$ and $\partial_t \sim \epsilon \partial_r$ for the geometric fields (see Seidel & Suen 1990; Guzmán & Ureña-López 2004).

Consistent with the $T_{\alpha\beta}$ computed in the flat space-time, the right hand side elements in Einstein's equations are written as

$$\begin{aligned} S_{tt} &= \Phi_{,t}\Phi_{,t} - \frac{1}{2}m^2|\Phi|^2, \\ S_{rr} &= \Phi_{,r}\Phi_{,r} + \frac{1}{2}m^2|\Phi|^2, \\ S_{tr} &= \Phi_{,r}\Phi_{,t}. \end{aligned} \quad (24)$$

In this case it is convenient to introduce the dimensionless quantities

$$x = mr, \quad \tau = mt, \quad \Omega = \frac{\omega}{m}, \quad (25)$$

where we note that the bosonic mass m is the natural scale for time and distance. In terms of these new variables the general solution to equation (13) takes the form

$$\Phi(\tau, x) = \frac{1}{x} \exp(\pm ix\sqrt{\Omega^2 - 1}) \exp(\pm i\Omega\tau). \quad (26)$$

The physical properties of the solution depend on the ratio $\Omega \equiv \omega/m$. For $\Omega < 1$ the solution decays exponentially but it is singular at $r = 0$. On the other hand, $\Omega > 1$ allows for non-singular solutions which vanish at infinity. We will restrict ourselves to this case. We will write the particular solution in the form

$$\sqrt{\kappa_0}\Phi(\tau, x) = \phi(x) \cos(\Omega\tau), \quad (27)$$

where the spatial function is given by

$$\phi(x) = \phi_0 \frac{\sin(x\sqrt{\Omega^2 - 1})}{x}. \quad (28)$$

Because of the functional form of the scalar field (equation 27) we introduce the following *ansatz* for the metric perturbations (equation 21)

$$\begin{aligned} V(\tau, x) &= V_0(x) + V_2(x) \cos(2\Omega\tau), \\ U(\tau, x) &= U_0(x) + U_2(x) \cos(2\Omega\tau). \end{aligned} \quad (29)$$

We adopt the following method to solve the EKG equations. In the first approximation we substitute the solution of a lower approximation (the flat case) into the next approximation (with ϵ^2) and solve the resulting differential equations. As we will see, this standard approximation works well in our case. In terms of these expressions the Einstein's equations $R_{\alpha\beta} = \kappa_0 S_{\alpha\beta}$ finally read

$$\begin{aligned} \epsilon^2 \left[\frac{4}{x} U_2 \right] &= \frac{1}{2} \phi \phi_{,x}, \quad (30) \\ \epsilon^2 \left[V_{0,xx} + \frac{2}{x} V_{0,x} \right] &= \frac{1}{2} \left(\Omega^2 - \frac{1}{2} \right) \phi^2, \\ \epsilon^2 \left[V_{2,xx} + \frac{2}{x} V_{2,x} + 4\Omega^2 U_2 \right] &= -\frac{1}{2} \left(\Omega^2 + \frac{1}{2} \right) \phi^2, \\ \epsilon^2 \left[-V_{0,xx} + \frac{2}{x} U_{0,x} \right] &= \frac{1}{2} \left(\phi_{,x}^2 + \frac{1}{2} \phi^2 \right), \\ \epsilon^2 \left[-V_{2,xx} + \frac{2}{x} U_{2,x} - 4\Omega^2 U_2 \right] &= \frac{1}{2} \left(\phi_{,x}^2 + \frac{1}{2} \phi^2 \right). \end{aligned}$$

3.1. Scaling properties

From System (30) we know that the scalar field's maximum amplitude $\phi(0) = \phi_0 \sqrt{\Omega^2 - 1}$ could be taken as the expansion parameter ϵ and in this case Ω must be of order 1. Then it is always possible to solve the System (30) ignoring ϵ and replacing ϕ by its normalized function

$$\hat{\phi}(x) = \frac{\sin(x\sqrt{\Omega^2 - 1})}{x\sqrt{\Omega^2 - 1}}. \quad (31)$$

Solutions $\hat{\phi}$, U_0 , U_2 , V_0 , V_2 of this normalized system depend only of the arbitrary characteristic frequency

Ω which modulates the wave length of $\hat{\phi}$. On the other hand, for each value of Ω there is a complete family of solutions of the scalar field ϕ and the metric perturbations $h_{\alpha\beta}$ which are related to each other by a scaling transformation characterized by ϕ_0

$$\begin{aligned} \sqrt{\kappa_0}\Phi &= \phi_0 \sqrt{\Omega^2 - 1} \hat{\phi} \cos(\Omega\tau), \quad (32) \\ h_{rr} &= \phi_0^2 (\Omega^2 - 1) [2U_0 + 2U_2 \cos(2\Omega\tau)], \\ h_{tt} &= -\phi_0^2 (\Omega^2 - 1) [2V_0 + 2V_2 \cos(2\Omega\tau)], \end{aligned}$$

In this context the weak field limit condition $h_{\alpha\beta} \ll 1$ translates into

$$\phi_0^2 (\Omega^2 - 1) |2V| \ll 1, \quad \phi_0^2 (\Omega^2 - 1) |2U| \ll 1. \quad (33)$$

Here we will introduce a specific notation for the spatial functions of the metric perturbations:

$$\begin{aligned} h_{rr}^{(0)} &= \phi_0^2 (\Omega^2 - 1) 2U_0, \quad (34) \\ h_{rr}^{(2)} &= \phi_0^2 (\Omega^2 - 1) 2U_2, \\ h_{tt}^{(0)} &= \phi_0^2 (\Omega^2 - 1) 2V_0, \\ h_{tt}^{(2)} &= \phi_0^2 (\Omega^2 - 1) 2V_2. \end{aligned}$$

3.2. Metric perturbations solutions

The system of equations (30) can be solved and the spatial functions of the metric perturbations have analytic solutions given by

$$\begin{aligned} U_2 &= \frac{1}{8(\Omega^2 - 1)} \left[-\frac{\sin^2(x\sqrt{\Omega^2 - 1})}{x^2} + \frac{\sqrt{\Omega^2 - 1} \sin(2x\sqrt{\Omega^2 - 1})}{2x} \right], \\ V_0 &= \frac{(2\Omega^2 - 1)}{8(\Omega^2 - 1)} \left[\frac{\sin(2x\sqrt{\Omega^2 - 1})}{2x\sqrt{\Omega^2 - 1}} - Ci(2x\sqrt{\Omega^2 - 1}) + \ln(2x\sqrt{\Omega^2 - 1}) \right] - \frac{C_{V01}}{x} + C_{V02}, \\ V_2 &= \frac{1}{8(\Omega^2 - 1)} \left[\frac{\sqrt{\Omega^2 - 1} \sin(2x\sqrt{\Omega^2 - 1})}{2x} + Ci(2x\sqrt{\Omega^2 - 1}) - \ln(2x\sqrt{\Omega^2 - 1}) \right] + C_{V22}, \\ U_0 &= \frac{1}{8(\Omega^2 - 1)} \left[-\frac{1}{2x^2} + \frac{1}{2} \frac{\cos(2x\sqrt{\Omega^2 - 1})}{x^2} - \frac{1}{2} \frac{\sin(2x\sqrt{\Omega^2 - 1})}{x\sqrt{\Omega^2 - 1}} \right] + \frac{C_{V01}}{x} + C_{U01}, \end{aligned} \quad (35)$$

where Ci is the cosine integral function and C_{V01} , C_{V02} , C_{V22} , and C_{U01} are integration constants.

3.3. Weak Field Validity Range

From equation (13) it is evident that, in the limit in which we are working, the KG equation is decoupled from Einstein equations. Imposing regularity at the origin and asymptotic flatness to the KG solution we have chosen equation (27) with equation (28) as our scalar field particular solution where ϕ_0 and $\Omega > 1$ are still free parameters.

On the other hand, regularity at $x = 0$ requires $h_{rr}(x = 0, \tau) = 0$ which implies

$$C_{V01} = 0, \quad C_{U01} = \frac{\Omega^2}{8(\Omega^2 - 1)}; \quad (36)$$

then for the perturbations C_{V02} and C_{V22} are still free integration constants. Now we will describe the asymptotic behavior of these perturbations. Due to U_2 being at least one order of magnitude smaller than U_0 and its value oscillating around zero, it is U_0 which determines the behavior of h_{rr} . The U_0 value starts to oscillate, very near to the origin, around C_{U01} maintaining this behavior asymptotically. Then the asymptotic value of h_{rr} is the finite C_{U01} value. Contrary to this h_{tt} , due to the logarithm terms in V_0 and V_2 , h_{tt} is singular at infinity. Thus, the weak field condition (equation 33) is fulfilled only in a finite spatial region around the origin, *i.e.*, due the the approximation the solution is contained in a box, for which the walls are sufficiently far away from the center of the solution. We will say that this is the region where our weak field approximation is valid.

Unique solutions for the EKG system will be obtained fixing the ϕ_0 and Ω parameters and the constants C_{V02} and C_{V01} within the validity range of the approximation. As it is known, the measurement of potentials does not have physical sense by itself; then unique solutions will be determined through metric dependent observable quantities. Using the expressions given by equations (34), we can obtain the perturbed metric functions in terms of these solutions. In Figure 3 we present a plot of these metric perturbation functions, as well as of the scalar field, for two values of Ω .

It is important to mention that the width of the validity region where equation (33) is fulfilled, depends on Ω and ϕ_0 . This is because it is the factor ϕ_0^2 which modulate the perturbations, (see equation (32)). What it is not evident until the solutions (equation 35) are observed is that the Ω value, independently from ϕ_0 , could make the validity range width bigger. This is because in the logarithm argument there is the expression $\sqrt{\Omega^2 - 1}$, then as Ω is closer to one the logarithm terms rise more slowly.




Fig. 3. Profiles of $\phi(x)$, $h_{rr}^{(2)}(x)$, $h_{rr}^{(0)}(x)$, $h_{tt}^{(2)}(x)$ and $h_{tt}^{(0)}(x)$ with $\phi_0 = 1$; see text for details.

The order of magnitude for the other parameter in the metric perturbations, ϕ_0 , can be naturally determined from the asymptotic value taken by h_{rr} , which is reached very close to the origin

$$\lim_{x \rightarrow \infty} |h_{rr}| = \frac{\phi_0^2}{4} \Omega^2. \quad (37)$$

As Ω is nearly 1 the magnitude of h_{rr} is given by ϕ_0 . It is well known that for weak field systems like our Solar System the metric perturbations go like $h_{\alpha\beta} \sim 10^{-6}$. This value restricts our ϕ_0 to be $\phi_0 \lesssim 10^{-3}$.

3.4. Analytical Solutions vs Numerical Solutions

Analytical solutions ϕ_0 , $h_{rr}^{(2)}(x)$, $h_{rr}^{(0)}(x)$, $h_{tt}^{(2)}(x)$ and $h_{tt}^{(0)}(x)$ are shown in Figure 3. The value of ϕ_0 is 1 in both plots. As already noted, the value of Ω characterizes each family of solutions. Mainly Ω determines the wave length of ϕ and the rate of increase of $h_{tt}^{(0)}$ and $h_{tt}^{(2)}$; as Ω is closer to 1, this rate is smaller. These characteristics are shown in Figure 3.




Fig. 4. (Top) Spatial function ϕ of the scalar field Φ . Solid and dashed lines are analytical solutions with $\phi_0 = 0.001/(\Omega^2 - 1)^{1/2} = 0.001591$, and $\phi_0 = 0.002/(\Omega^2 - 1)^{1/2} = 0.001946$, respectively. The crosses are the corresponding numerical solutions with $\phi_1(0) = 0.001$, $\Omega = 1.181008$ and $\phi_1(0) = 0.002$, $\Omega = 1.433822$; see text for details. (Bottom) Spatial functions $h_{tt}^{(0)}$ and $h_{tt}^{(2)}$ of the metric perturbations.

The exact EKG equations for spherical symmetry with a quadratic potential were solved numerically in Ureña-López (2002) and Ureña-López et al. (2002), where the so-called oscillatons were found. In those works boundary conditions are determined by requiring non-singular and asymptotically flat solutions, for which the EKG become an eigenvalue problem. The free eigenvalue is the scalar field's central value $\phi_1(x = 0)$ which labels the particular equilibrium configuration, and the fundamental frequency Ω is an output value. In those works it was also noted that weak gravity fields are produced by oscillatons with $\phi_1(x = 0) \ll 1$. In Figure 4 we compare some of these numerical solutions (NS) with the analytical solutions (AS) within a central region. The constant values of the AS are fixed to better fit the NS inside the weak field validity range. From these plots we




Fig. 5. (Top) Energy-momentum density profiles for two scalar field configurations, one with $\phi_0 = 1.1591 \times 10^{-3}$ and $\Omega = 1.181008$, and the other with $\phi_0 = 5.2225 \times 10^{-3}$ and $\Omega = 1.153246$. (Bottom) Difference between the density profiles from the exact EKG equations (ρ_N) and the density computed in the flat space-time (ρ); see text for details.

can conclude that our solutions are a very good approximation to the exact EKG equations in the weak field limit. The principal advantage of this approximation is the analytical description of the solutions.

4. SCALAR FIELD AS DARK MATTER: HALO DENSITY PROFILE

In this section we explore whether or not the scalar field could account for the galactic DM halos. Specifically, we compare the SFDM model density profile and the profiles inferred throughout the rotation curves profiles of galaxies which are mostly formed by DM.

As long as we are concerned with perturbations of the flat space-time due to the scalar field, we do not consider the baryonic matter gravitational effects, we expect that our approximation will be better suited for galaxies with very small baryonic component.

TABLE 1
GALACTIC PARAMETER VALUES

Galaxy	Ω	ϕ_0	χ^2	$\rho(r=0)$ [M_\odot/pc^3]	$\Delta\rho(0)$ [M_\odot/pc^3]	T [yrs]
ESO0140040	$1+ 8 \times 10^{-9}$	1.87×10^{-3}	12.366	0.569×10^{-1}	0.755×10^{-8}	1.603382750×10^7
ESO0840411	$1+ 6 \times 10^{-9}$	5.95×10^{-4}	1.338	0.433×10^{-2}	0.147×10^{-9}	1.603953416×10^7
ESO1200211	$1+53 \times 10^{-9}$	2.04×10^{-4}	10.062	0.448×10^{-2}	0.530×10^{-9}	1.603382679×10^7
ESO1870510	$1+12 \times 10^{-9}$	3.28×10^{-4}	3.190	0.265×10^{-1}	0.699×10^{-8}	1.603382570×10^7
ESO2060140	$1+18 \times 10^{-9}$	9.18×10^{-4}	65.421	0.308×10^{-1}	0.206×10^{-8}	1.603382735×10^7
ESO3020120	$1+29 \times 10^{-9}$	6.88×10^{-4}	16.099	0.279×10^{-1}	0.170×10^{-8}	1.603382718×10^7
ESO3050090	$1+40 \times 10^{-9}$	4.72×10^{-4}	1.224	0.181×10^{-1}	0.243×10^{-8}	1.603382699×10^7
ESO4250180	$1+ 4 \times 10^{-9}$	1.27×10^{-3}	5.221	0.132×10^{-1}	0.105×10^{-8}	1.603382757×10^7
ESO4880049	$1+ 3 \times 10^{-9}$	7.86×10^{-4}	11.410	0.377×10^{-1}	0.212×10^{-8}	1.603382715×10^7




Fig. 6. Density profile fits for the galaxies ESO0140040, ESO2060140. The right panel is a zoom of the central region, plotted in order to visualize complete error bars. Units for the horizontal axes are kpc , for the vertical axes M_\odot/pc^3 . See text and Table 1 for fit details.

We will compare the energy-momentum density for the scalar field given by equation (7), in the relativistic weak limit approximation, where for the metric functions, equation (21), we use the solutions to the perturbations given by equations (29) and (35).

This is consistent with the fact that gravity does not modify the scalar field behavior. This approximation in the weak gravitational field limit is very good as we can see in Figure 5. In the plots we show the energy-momentum density profiles for two scalar field configurations with different maximum amplitudes at the origin $\phi(0) = \phi_0\sqrt{\Omega^2 - 1}$. It is

important to notice that as $\phi(0)$ decreases the gravitational field gets weaker, and the difference between the density from the complete EKG equations and from our approximation becomes smaller.

The density profile fits allow us to obtain an estimation of the parameters at the galactic level: the fundamental frequency Ω and the scalar field constant ϕ_0 . The third parameter involved in the density profiles is the scalar field mass; we will fix it to be $m = 10^{-23}$ eV. This value was fitted for the SFDM model from cosmological observations in Matos & Ureña-López (2001).




Fig. 7. The same as Figure 6, but for the galaxies ESO3020120, ESO3050090.




Fig. 8. The same as Figure 6, but for the galaxies ESO4250180, ESO1200211.

4.1. Density Profile Fits

The first qualitative feature of the energy-density profile that we want to emphasize is that in the central region it is non-cuspy (see Figure 5). It is important to take into account that instead of density profiles, rotation curves are the direct observable for galaxies. Nevertheless, for galaxies dominated by DM, their rotation curves could model the DM density profile more trustfully. We choose a subset of galaxies from the set presented in McGaugh, Rubin, & de Blok (2001), the common characteristic for the

selected galaxies is that the luminous matter velocity contribution to the rotation curves is almost null.

With the scalar field mass m fixed, the profile fits were made for the Ω and ϕ_0 values with a good χ^2 statistic, see Table 1. In most of the cases the non-central observational data were the better fitted points; those data points also had smaller error bars. The density profile fits are depicted in Figures 6, 7, 8, 9, where we show several galaxies with the density computed from the observed rotational profile versus the density obtained with our SFDM description. In

some of them we were able to compare regions within less than 0.5 kpc.

In Table 1 the fundamental frequency Ω is listed for each galaxy. We found that the temporal dependence for the energy-momentum density profile is harmonic with a temporal period $T = \pi/\Omega$. The column $\Delta\rho(0)$ corresponds to the maximum change in the central density for a period of time T . Finally for all the galaxies the ϕ_0 value lies well inside of the weak gravitation field limit $\phi_0 \lesssim 10^{-3}$.

5. CONCLUSIONS AND FUTURE PROSPECTS

We have found analytic solutions for the EKG equations, for the case when the scalar field is considered as a test field in a Minkowski background, and in the relativistic weak gravitational field limit to first order in the metric perturbations. With these solutions we have shown that non-trivial local behavior of the scalar field holds the collapse of an object formed from scalar field matter. The scalar field contains non trivial, natural effective pressures which stop the collapse and prevent the centers of these objects from having cusp-like density profiles. Even within this simple approximation it has been possible to fit, with relative success, the density profiles for some galaxies showing non-cuspy profiles.

Together, all the features of the SFDM model allow one to consider this model as a robust alternative candidate to be the dark matter of the Universe, as was suggested by Guzmán & Matos (2000), Matos & Guzmán (2000, 2001), and Matos, Guzmán, & Núñez (2000). Furthermore, it has been shown previously that dark halos of galaxies could be scalar solitonic objects, even in the presence of baryonic matter (Hu, Barkana, & Gruzinov 2000; Lee & Koh 1996; Arbey, Lesgourgues, & Salati 2001, 2002; Sin 1994; Ji & Sin 1994). Actually, the boson mass estimated in all these different approaches roughly coincides with the value $m_\phi \sim 10^{-23}\text{eV}$, even if the later was estimated from a cosmological point of view, Matos & Ureña-López (2001). We can appreciate the non-trivial characteristics of the proposed potential (equation 2): Its strong self-interaction provides a reliable cosmological scenario, while at the same time it has the desired properties of a quadratic potential. Finally, the results presented here fill the gap between the successes at cosmological and galactic levels.

We would like to thank Miguel Alcubierre, Vladimir Ávila Reese, Arturo Ureña and F. Siddhartha Guzmán for many helpful and useful discussions, and Erasmo Gómez and Aurelio Espíritu for




Fig. 9. Density profile fits for the galaxies ESO1870510, ESO488049, ESO0840411.

technical support. We thank Lidia Rojas for careful reading and suggestions to improve our work. The numeric computations were carried out in the “Laboratorio de Súper-Cómputo Astrofísico (LaSumA) del Cinvestav”. This work was partly supported by Conacyt México, under grants 32138-E, 47209-F and 42748 and by grant number I0101/131/07 C-234/07, Instituto Avanzado de Cosmología (IAC) collaboration. Also by the bilateral Mexican-German project DFG-Conacyt 444 MEX-13/17/0-1.

REFERENCES

- Alcubierre, M., Becerril, R., Guzmán, F. S., Matos, T., Núñez, D., & Ureña-López, L. A. 2003, *Classical Quantum Gravity*, 20, 2883
- Alcubierre, M., Guzmán, F. S., Matos, T., Núñez, D., Ureña-López, L. A., & Wiederhold, P. 2002, *Classical Quantum Gravity*, 19, 5017
- Arbey, A., Lesgourgues, J., & Salati, P. 2001, *Phys. Rev. D*, 64, 123528
- _____. 2002, *Phys. Rev. D*, 65, 083514
- Böhmer, C. G., & Harko, T., 2007, *J. Cosmol. Astropart. Phys.*, 6, 25
- Guzmán, F. S., & Matos, T. 2000, *Classical Quantum Gravity*, 17, L9
- Guzmán, F. S., & Ureña-López, L. A. 2004, *Phys. Rev. D*, 69, 124033
- _____. 2005, in *Progress in Dark Matter*, ed. J. Val Blain (New York: Nova Science), 39
- Hawley, S. H., & Choptuik, M. W. 2000, *Phys. Rev. D*, 62, 104024
- Honda, E. P., & Choptuik, M. W. 2002, *Phys. Re. D*, 65, 084037
- Hu, W., Barkana, R., & Gruzinov, A. 2000, *Phys. Rev. Lett.*, 85, 1158
- Ji, S. U., & Sin, S. J. 1994, *Phys. Rev. D*, 50, 3655
- Lee, J. W., & Koh, I. G. 1996, *Phys. Rev. D*, 53, 2236
- MacGaugh, S. S., Rubin, V. C., & de Blok, W. J. G. 2001, *AJ*, 122, 2381
- Matos, T., & Guzmán, F. S. 2000, *Ann. Phys.*, 9, S133
- _____. 2001, *Classical Quantum Gravity*, 18, 5055
- Matos, T., Guzmán, F. S., & Núñez, D. 2000, *Phys. Rev. D*, 62, 061301
- Matos, T., & Ureña-López, L. A. 2000, *Classical Quantum Gravity*, 17, L75
- _____. 2001, *Phys. Rev. D*, 63, 063506
- Navarro, J., Frenk, C. S., & White, S. S. M. 1997, *ApJ*, 490, 493
- Peebles, P. J. E. 2000, *ApJ*, 534, L127
- Sahni, V., & Wang, L. 2000, *Phys. Rev. D*, 62, 103517
- Schunck, F. E. 1998, preprint (astro-ph/9802258)
- Seidel, E., & Suen, W.-M. 1990, *Phys. Rev. D*, 42, 384
- _____. 1991, *Phys. Rev. Lett.*, 66, 1659
- _____. 1994, *Phys. Rev. Lett.*, 72, 2516
- Sin, S. J. 1994, *Phys. Rev. D*, 50, 3650
- Ureña-López, L. A. 2002, *Classical Quantum Gravity*, 19, 2617
- Ureña-López, L. A., Matos, T., & Becerril, R. 2002, *Classical Quantum Gravity*, 9, 6259

Argelia Bernal and Tonatiuh Matos: Departamento de Física, Centro de Investigación y de Estudios Avanzados del IPN, Apdo. Postal 14-740, 07000 México, D. F., Mexico <http://www.fis.cinvestav.mx/~tmatos> (abernal, tmatos@fis.cinvestav.mx).

Darío Núñez: Instituto de Ciencias Nucleares, Universidad Nacional Autónoma de México, Apdo. Postal 70-543, 04510 México, D. F., Mexico. On Sabbaticak leave at Max-Planck-Institut für Gravitationsphysik, Albert Einstein Institut, 14476 Golm, Germany (nunez@nuclecu.unam.mx).