

The species of Entiminae (Coleoptera: Curculionidae) ranged in America south of the United States

JUAN J. MORRONE*

Resumen. Se listan los nombres válidos de los géneros y especies de entiminos (Coleoptera: Curculionidae: Entiminae) distribuidos en América al sur de los Estados Unidos, incluyendo sus sinónimos y sumarios distribucionales. Estos taxones pertenecen a dos tribus: Sitonini y Entimini, esta última incluyendo las subtribus Leptopiina (= Cylydrorhinina, syn. nov., Strangalioidina, Rhigopsina y Premnotrypina, syn. nov.), Entimina (= Lordopina, syn. nov., Hypsonotina, syn. nov., Promecopina, syn. nov. y Eudiagogina, syn. nov.), Ophryastina, Eustylina (=Exophthalmina y Exophthalmodina), Otiorhynchina (= Oosomina, Peritelina, Trachyphloeina y Brachyrhinina), Polydrosina (= Brachyderina, Cneorhinina y Scythropina), Anypotactina, Barynotina, Anomophthalmina, Tanymecina (= Piazomiina y Pandeletiina) y Naupactina (= Cyphina, Neocyphina y Platymolina). Se establecen 52 nuevas sinonimias de nombres del grupo de la especie, y se da estatus específico a cuatro subespecies.

Palabras clave: Coleoptera, Curculionidae, Entiminae, América.

Abstract. The valid names of the genera and species of entimines (Coleoptera: Curculionidae: Entiminae) ranged in America south of the United States are listed, including their synonyms and distributional summaries. These taxa belong in two tribes: Sitonini and Entimini, the latter including the subtribes Leptopiina (= Cylydrorhinina, syn. nov., Strangalioidina, Rhigopsina, and Premnotrypina, syn. nov.), Entimina (= Lordopina, syn. nov., Hypsonotina, syn. nov., Promecopina, syn. nov., and Eudiagogina, syn. nov.), Ophryastina, Eustylina (=Exophthalmina and Exophthalmodina), Otiorhynchina (= Oosomina, Peritelina, Trachyphloeina, and Brachyrhinina), Polydrosina (= Brachyderina, Cneorhinina, and Scythropina), Anypotactina, Barynotina, Anomophthalmina, Tanymecina (= Piazomiina and Pandeletiina), and Naupactina (= Cyphina,

* Museo de Zoología, Facultad de Ciencias, Universidad Nacional Autónoma de México (UNAM), Apartado postal 70-399, 04510 México D.F., México.

Neocyphina, and Platynomina). A total of 52 new synonymies of species-group names are established, and four subspecies are given species status.

Key words: Coleoptera, Curculionidae, Entiminae, America.

Introduction

The subfamily Entiminae (Coleoptera: Curculionidae) comprises the weevil taxa formerly known collectively as Adelognatha, and includes approximately 1 150 genera and 12 220 species, distributed in all biogeographical regions of the world (Thompson 1992). Kuschel (1995) demoted Entiminae to a tribe of Brachycerinae, although the majority of authors (Thompson 1992, May 1994, Zimmerman 1994, Morris 1995, Morrone 1996, 1997, 1998, Marvaldi 1997, 1998) treat them as a distinct subfamily. The majority of adult Entiminae share adelognathous mouthparts and deciduous mandibular processes (Thompson 1992), but these characters have proved to be highly homoplastic and are also present in other weevil taxa (Marvaldi 1997). Two larval synapomorphies, however, justify the monophyly of this subfamily: the wider than long, dorsoventrally compressed, cushion-like antennal sensorium, and the maxillary mala with four ventral setae (Marvaldi 1997, 1998). According to a recent cladistic analysis based on larval features (Marvaldi 1997), they are classified into five tribes: Pachyrhynchini, Ectemnorhinini, Alohini, Sitonini, and Entimini. Two of these tribes, Sitonini and Entimini, are distributed in the New World.

Checklists providing basic information are helpful when working on taxonomic revisions and catalogs. O'Brien & Wibmer (1982) & Wibmer and O'Brien (1986) compiled the most recent checklists of the New World weevils. Over the course of the last years, however, several new entimine taxa have been described and new combinations and synonymies established subsequent to these checklists (O'Brien & Wibmer 1984, Howden 1986, 1993a, b, 1996, 1998, Tanner 1987, Wibmer & O'Brien 1989, Lanteri 1990a-d, 1992, 1995, Lanteri & O'Brien 1990, Bordón 1991, 1997, Lanteri & Morrone 1991, 1995, Bright 1994; Lanteri & Díaz 1994; Normark & Lanteri 1996; Morrone 1997, 1998).

My objective is to provide a list of the entimines ranged in America south of the United States, incorporating changes and additions that update O'Brien & Wibmer's (1982) and Wibmer & O'Brien's (1986) checklists.

For the subtribal names of Entimini I include herein only some selected synonyms, that are relevant for the New World taxa. The arrangement of the genera and species within each tribe and subtribe is alphabetical. Each name is followed by the original citation, with its synonyms listed chronologically between square brackets. Distributional data for each species are summarized by country; an asterisk denotes that it is also present in America north of Mexico.

The polytypic species concept, which has led to recognition of subspecies whenever a low degree of differentiation seemed to not justify a species status for allopatrid populations, has come under strong criticism and has been largely abandoned in modern systematics (Cracraft 1992). For this reason, I have chosen to synonymize the names of subspecies and varieties, unless they could be treated as distinct species.

Subfamily ENTIMINAE Schoenherr 1823
Tribe SITONINI LeConte 1874

Sitona Germar 1817: 341 [= *Sitones* Schoenherr 1840: 253, unjust. emend.]

- S. californicus* (Fahraeus 1840: 267) [= *S. ordinarius* (Fahraeus 1840: 266); *S. angustulus* (Casey 1888: 292); *S. apacheanus* (Casey 1888: 294); *S. explicitus* (Casey 1888: 293); *S. extrusus* (Casey 1888: 282); *S. hispidiceps* (Casey 1888: 292); *S. occidentalis* (Casey 1888: 285); *S. prominens* (Casey 1888: 291); *S. cockerelli* Blaisdell, 1938: 31; *S. bryanti* Tanner, 1987: 172; *S. oregonensis* Tanner, 1987: 172]. *Mexico.
S. cylindricollis (Fahraeus 1840: 269) [= *S. procerus* (Casey 1888: 284); *S. alpinensis* Tanner 1987: 170]. Asia and Europe. Introduced into the United States and Mexico.
S. discoideus Gyllenhal 1834: 112. Africa and Europe. Introduced into Australia, Chile, and New Zealand.
S. hispidulus (Fabricius 1776: 226). *Mexico.
S. lepidus Gyllenhal 1834: 104 [= *S. flavescentis* of authors, not Marsham 1801]. *Mexico.
S. lineellus (Bonsdorff 1785: 30) [= *S. indifferens* Say 1831: 10; *S. scissifrons* Say 1831: 10; *S. vittatus* (Couper 1865: 85)]. North America. Introduced into Mexico.
S. vittatus (LeConte 1857: 54) [= *S. sordidus* (LeConte 1857: 54); *S. alternans* (Casey 1888: 288); *S. eximus* (Casey 1888: 286); *S. margaritosus* (Casey 1888: 284); *S. montanus* (Casey 1888: 287); *S. osculans* (Casey 1888: 290)]. *Mexico.

Tribe ENTIMINI Schoenherr 1823
Subtribe LEPTOPIINA Lacordaire 1863

Cylydrorhinina Lacordaire 1863, **syn. nov.**

Strangalioidina Lacordaire 1863.

Rhigopsina LeConte 1874.

Premnotrypina Kuschel 1956, **syn. nov.**

Adaleres Casey 1895: 816

A. humeralis Casey 1895: 818. *Mexico.

Amphidees Schoenherr 1842: 252 [= *Deamphus* Sharp 1891: 102]

A. acuminatus Champion 1911: 322. Mexico.

A. alternans Sharp 1891: 99. Mexico.

A. brevipennis (Sharp 1891: 103). Guatemala.

A. deceptor (Sharp 1891: 103). Guatemala and Mexico.

A. latifrons (Sharp 1891: 103). Mexico.

A. longulus Sharp 1891: 100. Mexico.

A. macer Sharp 1891: 98. Mexico.

A. major Sharp 1891: 97. Mexico.

A. nasutus Sharp 1891: 99. Mexico.

A. nubilosus Boheman 1842: 253. Mexico.

A. puncticollis (Sharp 1891: 104). Mexico.

Bastactes Schoenherr 1842: 293

B. bituberculatus Boheman 1842: 294 [= *B. conifer* Redtenbacher 1867: 151]. Brazil.

B. plurituberculatus Dohrn 1884: 43. Brazil.

Bolivianus Hustache 1923: 292

B. variabilis Hustache 1923: 293. Argentina, Bolivia, and Paraguay.

Caneorhinus Kuschel 1952: 121

C. biangulatus (Champion 1918: 177). Argentina (Islas Malvinas).

C. gravidus (Burmeister 1879: 208). Argentina and Chile.

C. lineatus (Blanchard 1851: 355) [= *C. confusaneus* (Berg 1899: 61); *C. confuseanus* (Enderlein 1912: 79), lapsus]. Argentina and Chile.

C. tessellatus (Guérin 1839: 303). Argentina and Chile.

C. uretai Kuschel 1958: 251. Argentina and Chile.

Cimbocera Horn 1876: 55

C. pilosa (Sharp 1891: 100). Mexico.

Cindynus Kuschel 1955: 299

C. sericeus (Blanchard 1851: 378) [= *C. lateralis* (Blanchard 1851: 378)]. Chile.

Cylydrorhinus Guérin 1830: 119 [= *Otideres* Dejean 1835: 260, nom. nud.; *Adioristus* Waterhouse 1841: 124; *Cylindrorthinus* Schoenherr 1842: 336, lapsus; *Scotoeborus* Schoenherr 1843: 97; *Otideres* Fairmaire & Germain 1860: 4; *Cylindrorthinus* Lacordaire 1863: 341, lapsus; *Otidoderes* Lacordaire 1863: 343; *Cylindrorthinus* Gemminger & Harold 1871: 2358, lapsus; *Amathynetes* Olliff 1891: 70; *Nahuelius* Germain 1895: 313; *Paulsenius* Germain 1895: 318; *Nahuelis* Germain 1911: 211, lapsus; *Otidores* Germain 1911: 212, lapsus; *Otiodes* Bruch 1915: 413, lapsus; *Listroderes* Champion 1918: 178, misident., not Schoenherr 1826; *Peradioristus* Voss 1954: 242; *Protadioristus* Voss 1954: 241; *Pseudadioristus* Voss 1954: 243]

C. albostrigosus (Fairmaire & Germain 1860: 7). Chile.

C. alticola (Olliff 1891: 71). Ecuador.

C. angulatus Guérin 1841: 217 [= *C. rubripes* (Philippi 1862: 410); *C. rugatus* (Philippi 1862: 411); *C. scrobiculatus* (Burmeister 1879: 203)]. Argentina and Chile.

C. angustatus (Waterhouse 1841: 125) [= *C. magellanicus* (Blanchard 1853: 209)]. Chile.

C. araucanus Kuschel 1952: 121 [= *C. araucariae* (Germain 1911: 212), nom. nud.]. Chile.

C. aspericollis (Fairmaire 1884: 502) [= *C. aspericollis* (Bruch 1915: 413), lapsus; *C. carinirostris* (Voss 1932: 42)]. Argentina and Chile.

C. aureolus (Hustache 1926: 190). Argentina.

C. aymaranus (Kuschel 1949: 37). Bolivia, Chile, and Peru.

C. birabeni Kuschel 1958: 250. Argentina.

- C. boliviensis* (Hustache 1938: 179). Bolivia.
- C. brachyderoides* (Hustache 1926: 188). Argentina.
- C. bruchi* Kuschel 1958: 248. Argentina.
- C. burmeisteri* Kuschel 1958: 245 [= *C. patagonicus* Germain 1911: 212, *nom. nud.*]. Argentina and Chile.
- C. carinicollis* (Blanchard 1851: 340) [= *C. corralensis* (Germain 1895: 488)]. Chile.
- C. caudatus* (Voss 1954: 248). Peru.
- C. caudiculatus* (Fairmaire 1889: 131) [= *C. caudiculatus* (Fairmaire 1884: 503), *nom. nud.*; *C. carinicollis* Germain 1895: 487, misident., not Blanchard, 1851; *C. caudiculatus* (Germain 1895: 486); *C. medianus* (Schenkling & Marshall 1931: 6)]. Argentina (Islas Malvinas) and Chile.
- C. chilensis* (Blanchard 1851: 326) [= *C. montanus* (Lacordaire 1863: 342)]. Argentina and Chile.
- C. clathratus* Blanchard 1851: 356. Argentina and Chile.
- C. comatus* (Erichson 1847: 129). Peru.
- C. conspersus* (Waterhouse 1841: 125) [= *C. araucanus* (Germain 1911: 211, *nom. nud.*; *C. humilis* (Germain 1911: 212, *nom. nud.*; *C. acuminatus* (Hustache 1926: 191)]. Argentina and Chile.
- C. costatus* (Blanchard 1851: 352) [= *C. griseus* (Philippi 1862: 411); *C. apiculatus* (Hustache 1938: 281)]. Argentina and Chile.
- C. cupreosquamulosus* (Germain 1893: 57). Chile.
- C. denticulatus* (Fairmaire & Germain 1860: 4) [= *C. inquinatus* (Lacordaire 1863: 343)]. Chile.
- C. dentipennis* (Burmeister 1879: 205). Argentina and Chile.
- C. echinosoma* (Fairmaire 1884: 501). Argentina.
- C. elongatus* (Kuschel 1949: 41). Chile and Peru.
- C. farinosus* (Burmeister 1879: 207). Argentina and Chile.
- C. fortunai* (Bruch 1940: 113). Argentina.
- C. frigidus* (Germain 1895: 318). Argentina and Chile.
- C. fulvipes* (Guérin 1839: 304) [= *C. sulcatus* (Fairmaire 1885: 60); *C. fuegianus* (Berg 1899: 63)]. Argentina and Chile.
- C. gemignanii* Kuschel 1958: 243. Argentina.
- C. glaberrimus* Kuschel 1958: 242. Argentina and Chile.
- C. gonoderus* (Fairmaire & Germain 1860: 7). Chile.
- C. granulatus* Kuschel 1958: 249. Argentina.
- C. hercules* Kuschel 1958: 246. Argentina.
- C. herrerai* (Kuschel 1949: 29). Bolivia and Chile.
- C. hirtus* (Voss 1954: 249). Peru.
- C. horridus* (Burmeister 1879: 208). Argentina.
- C. inaequatus* (Schenkling & Marshall 1931: 8) [= *C. inaequalis* (Germain 1896: 737), not Blanchard 1851]. Chile.
- C. inconspicuus* (Olliff 1891: 69). Ecuador.
- C. indefensus* Kuschel 1952: 124. Argentina and Chile.
- C. lactifer* (Burmeister 1879: 206) [= *C. externevittatus* (Fairmaire 1884: 501); *C. quadrilineatus* (Bovie 1907: 70)]. Argentina and Chile.
- C. laevipennis* Kuschel 1958: 244. Argentina and Chile.
- C. lateralis* (Berg 1899: 64). Argentina and Chile.

- C. laticollis* (Kuschel 1952: 126). Argentina and Chile.
- C. lemniscatus* (Quoy & Gaimard 1824: 549) [= *C. arrogans* (Boheman 1843: 204); *C. bicaudatus* (Enderlein 1907: 47)]. Argentina (Islas Malvinas).
- C. leprus* (Hustache 1938: 284). Bolivia and Peru.
- C. lineatus* (Kuschel 1949: 38). Bolivia and Chile.
- C. lineiger* (Kuschel 1958: 239) [= *C. vagelineatus* (Germain 1911: 212, *nom. nud.*)]. Argentina and Chile.
- C. manni* (Kuschel 1949: 40). Bolivia and Chile.
- C. marinus* (Burmeister 1879: 204) [= *C. murinus* Kuschel 1958: 241, not Boheman 1843, unjust. emend.]. Argentina.
- C. melanoleucus* (Burmeister 1879: 207) [= *C. aysenensis* Kuschel 1958: 248; *C. chubutensis* Kuschel 1958: 247]. Argentina and Chile.
- C. metrius* Kuschel 1952: 123. Argentina and Chile.
- C. mucronatus* (Hustache 1926: 191). Argentina.
- C. murinus* (Boheman 1843: 98). Argentina.
- C. nahuelius* (Kuschel 1949: 33) [= *C. atacamensis* (Germain 1911: 211), *nom. nud.*]. Argentina, Bolivia, and Chile.
- C. obesus* (Burmeister 1879: 205). Argentina.
- C. oblongus* Blanchard 1851: 357 [= *C. crassirostris* (Fairmaire & Germain 1860: 7)]. Chile.
- C. parvulus* (Blanchard 1851: 349). Chile.
- C. patagonicus* (Kuschel 1958: 238). Argentina and Chile.
- C. percostatus* (Fairmaire & Germain 1860: 8) [= *C. costatus* Redtenbacher 1867: 154]. Chile.
- C. perforatus* (Fairmaire & Germain 1860: 7). Chile.
- C. pilifrons* (Kirsch 1889: 28). Ecuador.
- C. pilosissimus* (Schenkling & Marshall 1931: 9) [= *C. pilosus* (Boheman 1842: 190), not Waterhouse 1841]. Chile.
- C. pilosus* (Waterhouse 1841: 122) [= *C. sitonoides* (Blanchard 1851: 353); *C. trichophorus* (Philippi 1887: 140)]. Chile.
- C. platyderes* Kuschel 1986: 106 [= *C. planicollis* Kuschel 1958: 242, not Blanchard 1851]. Argentina.
- C. plumeus* (Kuschel 1949: 35). Bolivia, Chile, and Peru.
- C. praelongus* (Fairmaire & Germain 1860: 7). Chile.
- C. pubescens* (Hustache 1926: 187). Argentina.
- C. pumilus* (Hustache 1926: 192). Argentina.
- C. punctatissimus* (Olliff 1891: 69). Ecuador.
- C. punctulatus* (Waterhouse 1841: 124). Chile.
- C. quadricollis* (Kuschel 1958: 240) [= *C. lineatus* (Burmeister 1879: 204), misident., not Blanchard 1851]. Argentina and Chile.
- C. setosulus* (Voss 1954: 242). Chile.
- C. simplex* (Waterhouse 1841: 126). Chile.
- C. simulans* (Olliff 1891: 71). Ecuador.
- C. sordidus* (Burmeister 1879: 206) [= *C. schythei* (Germain 1895: 471)]. Argentina and Chile.
- C. squamirostris* (Germain 1896: 825). Chile.
- C. squamulatus* (Hustache 1926: 189). Argentina.
- C. subcostatus* (Waterhouse 1841: 121) [= *C. planicollis* (Blanchard 1851: 338); *C. thermarum* (Germain 1893: 56)]. Chile.

- C. subdenudatus* (Waterhouse 1841: 126). Argentina.
C. subsquameus (Hustache 1938: 282) [= *C. callanganus* (Voss 1954: 252)]. Peru.
C. titschacki (Voss 1954: 251). Peru.
C. ursinus (Germain 1896: 836). Chile.
C. vestitus (Voss 1954: 249). Peru.
C. villosulus (Hustache 1926: 189) [= *C. griseus* (Germain 1911: 211), *nom. nud.*; *C. unicolor* (Hustache 1926: 189)]. Argentina, Bolivia, and Chile.
C. vittatus (Guérin 1839: 304). Chile.
- Cyphometopus* Blanchard 1851: 314
C. angustus Blanchard 1851: 315. Chile.
C. argentatus (Blanchard 1851: 362). Chile.
C. aricensis Kuschel 1949: 27. Chile.
C. auratus (Blanchard 1851: 313) [= *C. fulgidivittatus* (Blanchard 1851: 314)]. Chile.
C. cinereus (Blanchard 1851: 363) [= *C. angustatus* (Blanchard 1851: 364); *C. cylindricollis* Boheman, 1859: 122]. Chile.
C. marmoratus (Blanchard 1851: 363). Chile.
C. masafuerae Aurivillius 1931: 464. Argentina and Chile.
C. minimus (Blanchard 1851: 317). Chile.
C. tessellatipennis Blanchard 1851: 315. Chile.
- Cnemotricha* Emden 1939: 425 [= *Cnemotrichia* Blackwelder 1947: 797, lapsus]
C. setifera (Boheman 1840: 917). Brazil.
- Dasydema* Blanchard 1851: 379
D. anucella Kuschel 1952: 240. Chile.
D. hirtella Blanchard 1851: 379. Argentina and Chile.
- Enthus* Schoenherr 1826: 141 [= *Rembus* Germar 1824: 378, not Latreille 1817; *Gastrodora* Berthold 1827: 385]
E. auricinctus (Germar 1824: 379). Brazil.
- Erepsimus* Schoenherr 1845: 408 [= *Eripsimus* Agassiz 1846: 142, unjust. emend.; *Eripsimus* Gemminger & Harold, 1871: 2357, unjust. emend.]
E. setifer Boheman 1845: 409 [= *E. setiger* Lacordaire 1863: 337, lapsus]. Argentina, Brazil, and Paraguay.
- Geniocremnus* Kuschel 1949: 20
G. angustirostris (Blanchard 1851: 368) [= *G. cognatus* (Blanchard 1851: 369)]. Chile.
G. chilensis (Boheman 1842: 269). Chile.
G. laticollis (Blanchard 1851: 370). Chile.
G. villosus (Blanchard 1851: 369). Chile.
- Geonemides* Blanchard 1851: 359
G. ater Blanchard 1851: 360. Chile.
- Heteractus* Kuschel 1955: 277
H. vagabundus (Boheman 1840: 455). Brazil.

Hustachius Marshall 1933: 62

H. americanus (Hustache 1926: 171). Argentina.

Hybreoleptops Kuschel 1949: 19

H. aureosignatus (Blanchard 1851: 367). Chile.

H. tuberculifer (Boheman 1842: 226) [= *H. margaritaceus* (Erichson 1847: 129); *H. tuberculifer* (Blanchard 1851: 368), not Boheman 1842]. Argentina and Chile.

H. vestitus (Blanchard 1851: 367). Argentina and Chile.

H. xanthomelas (Fairmaire & Germain 1861: 7). Chile.

Machaerophrys Kuschel 1958: 236

M. insuetus Kuschel 1958: 237. Argentina.

Malvinius Kuschel 1950: 11

M. compressiventris (Enderlein 1907: 58). Argentina (Islas Malvinas).

M. nordenskioeldi (Enderlein 1907: 48) [= *M. divaricatus* (Enderlein 1907: 46)]. Argentina (Islas Malvinas).

Megalometides Kuschel 1958: 778

M. cacicus Kuschel 1958: 779 [= *M. cacicus* Germain 1911: 218, nom. nud.]. Chile.

M. discors Kuschel 1958: 781. Chile.

Megalometes Schoenherr 1842: 267

M. andigena Kuschel 1952: 238 [= *M. angustirostris* Germain 1911: 218, misident., not Blanchard 1851]. Chile.

M. aureosquamoides (Boheman 1845: 428) [= *M. squamifer* Blanchard 1851: 366]. Chile.

M. spinifer Boheman 1842: 268. Argentina and Chile.

Micramphidees Wibmer & O'Brien 1986: 87 [= *Micramphidees* Voss 1954: 237, not available, type species not fixed]

M. castor Voss 1954: 238. Peru.

M. pollux Voss 1954: 239. Peru.

Microtrypes Kuschel 1956: 195

M. acutiventris Kuschel 1956: 197. Bolivia.

M. convexicollis Kuschel 1956: 200. Bolivia.

M. coriaceus Kuschel 1956: 198. Bolivia.

M. curvipes Kuschel 1956: 199. Bolivia.

M. dissimilis (Hustache 1938: 182). Bolivia.

M. illimaniensis Kuschel 1956: 202. Bolivia.

M. pacis Kuschel 1956: 201. Bolivia.

M. puberulus Kuschel 1956: 219. Peru.

M. rotundicollis Kuschel 1956: 198. Bolivia.

M. striatus Kuschel 1956: 200. Bolivia.

M. tunariensis Kuschel 1956: 202. Bolivia.

Miostictus Pascoe 1886: CLIV

M. volxemi Pascoe 1886: CLIV. Brazil.

Neohustachius Hustache 1940: 112

N. tuberculipennis Hustache 1940: 112. Argentina.

Opseotapinotus Heller 1921: 30 [= *Opsiotapinotus* Kuschel 1949: 19, lapsus]

O. bicolor Hustache 1926: 164. Argentina.

O. caudatus Hustache 1926: 164. Argentina.

O. crinitus Heller 1921: 30. Argentina.

O. dicomus Kuschel 1945: 132. Argentina.

O. farinosus Kuschel 1945: 135. Argentina.

O. fricabilis Kuschel 1945: 133. Argentina.

O. molitor Heller 1921: 31. Argentina.

Orimodema Horn 1876: 43

O. protracta Horn 1876: 44. *Mexico.

Pantomorops Kuschel 1949: 20

P. striatus (Hustache 1947: 133). Argentina.

Paranametis Burke 1960: 123

P. distincta Burke 1960: 124. *Mexico.

Parergus Kuschel 1955: 300

P. albescens (Blanchard 1851: 345). Chile.

P. angusticeps (Blanchard 1851: 378). Chile.

P. axillaris (Fairmaire & Germain 1861: 7). Chile.

P. fascioliger (Blanchard 1851: 345). Chile.

Polycomus Schoenherr 1833: 653

P. lanuginosus Gyllenhal 1833: 654. Argentina and Brazil.

Premnotrypes Pierce 1914: 348 [= *Trypopremnon* Pierce 1914: 349; *Solanophagus* Hustache 1933: 376; *Plastoleptops* Heller, 1935: 276]

P. clivosus Kuschel 1956: 213. Bolivia.

P. fractirostris Marshall 1936: 497. Peru.

P. latithorax (Pierce 1914: 350). Bolivia, Chile, and Peru.

P. piercei Alcalá 1979: 63. Peru.

P. pusillus Kuschel 1956: 217. Peru.

P. sanfordi (Pierce 1918: 603). Peru.

P. solani Pierce 1914: 348. Peru.

P. solaniperra Kuschel 1956: 211. Bolivia and Peru.

P. solanivorax (Heller 1935: 276). Peru.

P. suturicallus Kuschel 1956: 215. Peru.

P. vorax (Hustache 1933: 377). Colombia, Ecuador, and Venezuela.

P. zischkai Kuschel 1956: 216. Bolivia.

Priocnemus Kirsch 1874: 391 [= *Priocnemis* Marshall 1944: 73, lapsus]

P. placidus (Erichson 1847: 129) [= *P. abendrothii* Kirsch 1874: 393]. Bolivia and Peru.

Rhigopsis LeConte 1874: 453

R. effracta LeConte 1874: 459 [= *R. scutellata* Casey 1888: 242]. *Mexico.

R. simplex Horn 1894: 442. Mexico.

Rhinotrypes Kuschel 1956: 189

R. grossepunctatus Kuschel 1956: 194. Bolivia and Peru.

R. juninensis Kuschel 1956: 192. Peru.

R. laevigatus Kuschel 1956: 191. Bolivia and Peru.

R. sulcifrons Kuschel 1956: 193. Ecuador.

R. weyrauchi Kuschel 1956: 219. Peru.

Sclerococcus Schoenherr 1840: 439

S. granulatus Fahraeus 1840: 440. Brazil.

Strangaliodes Schoenherr 1842: 219 [= *Malonotus* Blanchard 1851: 357; *Tapinopis* Blanchard

1851: 377, lapsus; *Tapinopsis* Blanchard 1851: 376; *Opseotapinus* Gemminger 1871: 2309;

Opseotropis Gemminger 1871: 119; *Strangaloides* Bruch 1915: 411, lapsus; *Chersostrangaliodes* Kuschel 1949: 20]

S. albosquamulosus Boheman 1842: 220 [= *S. albicans* (Germain 1911: 218), nom. nud.; *S. roseus* (Germain 1911: 218), nom. nud.]. Chile.

S. azurescens Kuschel 1949: 24 [= *S. tarapacanus* Kuschel 1949: 25]. Chile and Peru.

S. deserticola Kuschel 1949: 26. Chile.

S. elongatus Blanchard 1851: 364. Chile.

S. mutuarius Kuschel 1952: 239 [= *S. laevirostris* (Germain 1911: 218), nom. nud.; *S. arvalis* Kuschel 1952: 240; *S. ruralis* Kuschel 1952: 239]. Argentina and Chile.

S. niger (Blanchard 1851: 358). Chile.

S. squamiger (Voss 1947: 48). Chile.

S. sticticus Blanchard 1851: 362. Chile.

S. sulcatulus (Blanchard 1851: 377) [= *S. phalaeratus* (Blanchard 1851: 377), incorrect original spelling; *S. phaleratus* (Blanchard 1851: 377)]. Chile.

Synosomus Jekel 1858: 358

S. geonomoides Jekel 1858: 359 [= *S. geonomoides* Gemminger & Harold 1871: 2310, lapsus]. Central America.

Tetraphysus Lacordaire 1863: 248 [= *Dysphiles* Pascoe 1886: 416; *Probastactes* Brèthes 1910: 208]

T. ferrugatus (Pascoe 1886: 417) [= *T. quadrispinosus* (Brèthes 1910: 209); *T. serietuberculatus* (Hustache 1938: 280)]. Argentina, Bolivia, and Brazil.

Vossius Kuschel 1955: 301

V. albicinctus (Gyllenhal 1834: 160). Brazil.

V. cimetarius (Voss 1953: 427). Brazil.

V. dolosus (Pascoe 1886: CLIV). Brazil.

V. hispidulus (Gyllenhal 1834: 161). Brazil.

V. humilis (Hustache 1926: 168). Argentina and Paraguay.

V. jucundus (Voss 1934: 72). Brazil.

- V. longicollis* (Hustache 1926: 169). Brazil.
V. nebulosus (Gyllenhal 1834: 161). Brazil.
V. peregrinatus (Fahraeus 1840: 438). Brazil.
V. perplexus (Hustache 1926: 167). Argentina and Brazil.
V. plaumanni (Voss 1953: 426). Brazil.
V. robsi (Voss 1951: 368). Brazil.
V. signatus (Voss 1934: 74). Brazil.
V. viridifasciatus (Voss 1934: 73). Brazil.

Subtribe ENTIMINA Schoenherr 1823

Lordopina Schoenherr 1823, **syn. nov.**

Hypsonotina Jekel 1853, **syn. nov.**

Promecopina Lacordaire 1863, **syn. nov.**

Eudiagogina LeConte 1874, **syn. nov.**

Acanthobrachis Jekel 1854: I.9 bis [= *Acanthobrachys* Lacordaire 1863: 261, lapsus]
A. germari Jekel 1857: 103. Brazil and Colombia.

Aetherhinus Schoenherr 1840: 204 [= *Aetheorhinus* Agassiz 1846: 10; *Aetherrhinus* Gemminger and Harold 1871: 2226, lapsus]
A. aurichalceus Fahraeus 1840: 205. Brazil.

Aeuryomus Kessel 1935: 97

A. dubiosus Kessel 1935: 97. Brazil.

Alocorhinus Jekel 1854: I.6 [= *Alocorhinus* Sahlberg 1823: 32, not available, type species not fixed; *Alocorrhinus* Gemminger & Harold 1871: 2319, lapsus; *Alacorhinus* Faust 1892: 29, lapsus; *Allocorrhinus* Taschenberg 1907: 226, lapsus]

A. albolineatus (Sahlberg 1842: 146). Brazil.

A. comprimatus Sahlberg 1823: 32 [= *A. cathetoplateus* (Perty 1832: 74)]. Brazil.

A. curtisi (Sahlberg 1842: 148). Brazil.

A. gemmatus Jekel 1854: VI.15 [= *A. virescens* Jekel, 1857: 97; *A. squamulatus* Chevrolat 1881: CXLVI, nom. nud.]. Brazil.

A. hieroglyphicus Lucas 1857: 163. Brazil.

A. politus (Fahraeus 1842: 145). Brazil.

A. scrobiculatus (Fahraeus 1842: 147). Brazil.

A. succinctus (Fabricius 1776: 227) [= *A. albator* (Pallas 1781: 36); *A. marginellus* (Olivier 1790: 542), misident., not Fabricius 1781]. Brazil.

A. tener Kessel 1935: 90. Costa Rica.

Anidopsis Kessel 1935: 79 [not available, type species not fixed]

A. gracilior Kessel 1935: 81. Brazil.

A. molesta Kessel 1935: 81. Brazil.

A. perlata Kessel 1935: 80. Brazil.

Aracanthus Say 1831: 9 [= *Hyperacanthus* Voss 1934: 97, not available, type species not fixed; *Hyperacanthus* Blackwelder 1947: 806, lapsus]

- A. mourei* Rosado-Neto 1981: 69. Brazil.
A. notatus Voss 1934: 99. Brazil and French Guiana.
A. ovalis Voss 1934: 99. Brazil.
A. paraguayanus Voss 1943: 225 [= *A. paraguayanus* Voss 1943: fig. 1, lapsus]. Paraguay.
A. robustus Voss 1934: 98. French Guiana.
A. signatus Voss 1934: 100. Bolivia and Brazil.
A. tabaci Voss 1934: 98. Brazil.

Atomorhinus Hustache 1946: 3
A. impressidorsum Hustache 1946: 4. Colombia.

Aulametopus Jekel 1854: VI.7 [= *Aulacometopus* Gemminger & Harold 1871: 2319, unjust. emend.]
A. marginellus (Bohemian 1834: 256). Brazil.
A. marmoratus (Bohemian 1834: 255) [= *A. biacutus* (Fahraeus 1842: 177)]. Brazil.

Chileudius Kuschel 1950: 12
C. varians (Blanchard 1851: 324). Argentina and Chile.

Colecerus Schoenherr 1840: 927 [= *Colecerus* Agassiz 1846: 93, unjust. emend.; *Coleocerus* Gemminger & Harold 1871: 2374, unjust. emend.; *Bathyris* LeConte 1874: 461]
C. albidus (Chevrolat 1881: XXXVIII). Colombia.
C. crassipes (Champion 1911: 308). Mexico.
C. denticollis (Champion 1911: 307). Mexico.
C. dispar (LeConte 1874: 462). *Mexico.
C. marmoratus (Horn 1876: 109). *Mexico.
C. rotundicollis (Champion 1911: 309). Mexico.
C. setosus Boheman 1840: 928. Mexico.
C. variegatus Boheman 1845: 405. Costa Rica, El Salvador, Guatemala, Honduras, Mexico, and Nicaragua.
C. virescens (Champion 1911: 307). Mexico.

Conidus Marshall 1939: 581 [not available, type species not fixed] [= *Conus* Kessel 1935: 93, not Linnaeus 1758; *Conilla* Strand 1943: 96]
C. loculosus (Germar 1824: 370) [= *C. lacunosus* (Schoenherr 1834: 254), lapsus]. Brazil.
C. repertus (Kessel 1935: 95). Brazil.

Conothorax Jekel 1854: I.9 bis [= *Conothorax* Kessel 1935: 112, not Jekel 1854; *Conocloeus* Marshall 1939: 581]
C. albonotatus Bondar 1949: 179. Brazil.
C. luctuosus Lacordaire 1863: 277 [= *C. variabilis* Kessel 1935: 112]. Bolivia and Brazil.

Cydianerus Schoenherr 1840: 737 [= *Cydianirus* Agassiz 1846: 111, unjust. emend.; *Cydianirus* Gemminger & Harold 1871: 2326, unjust. emend.; *Dacnirus* Pascoe 1881: 300]
C. araneiformis (Dalman 1833: 450) [= *C. tumidus* (Dalman, 1833: 450)]. Brazil.
C. argenteus Boheman 1840: 741. El Salvador, Honduras, and Mexico.
C. bifasciatus Hustache 1926: 167 [= *C. collaris* (Voss 1934: 72), **syn. nov.**; *C. interruptus* (Voss 1934: 72), **syn. nov.**]. Argentina, Brazil, and Paraguay.

- C. bohemani* Boheman 1840: 738 [= *C. boehmani* Boheman 1840: 738, lapsus]. Brazil.
C. donceli (Bovie 1908: 4). Costa Rica.
C. flexuosus (Pascoe 1881: 301). Brazil.
C. latruncularius (Perty 1832: 70) [= *C. mannerheimii* (Gyllenhal 1833: 450); *C. wahlbergi* Boheman 1840: 740; *C. chevrolati* Blanchard 1847: 201, *syn. nov.*; *C. latruncularis* Vaurie 1952: 163, lapsus]. Argentina, Bolivia, Brazil, and Paraguay.
C. ornatus (Pascoe 1881: 300). Brazil.
C. pascoei (Bovie 1908: 43). Colombia and Panama.
C. virescens Lucas 1857: 150. Brazil.
C. wagneri Bovie 1907: 327. Venezuela.

Deroconus Jekel 1854: I.9 bis [= *Deraconus* Kessel 1932: 66, lapsus]
D. rufipes (Lacordaire 1863: 269). Colombia and Venezuela.

- Diaprosomus* Jekel 1854: I.7
D. bivittatus (Perroud 1853: 512). Brazil.
D. collaris Kessel 1932: 45. Brazil.
D. imperialis Kessel 1932: 43. Brazil.
D. magnificus (Sahlberg 1842: 167). Brazil.
D. regalis (Perroud 1853: 508). Brazil and Paraguay.
D. splendidus Kessel 1932: 47. Brazil.
D. suturalis Kessel 1932: 48. Brazil.
D. viridilineatus (Fahraeus 1842: 168). Brazil.

Dioryhinus Jekel 1854: I.9 bis [= *Diorirhinus* Lacordaire 1863 : 268, lapsus]
D. scutellumalbum Kessel 1932: 49. Brazil.

- Elytroxys* Jekel 1854: I.9 bis
E. inflatus Lacordaire 1863: 275 [= *E. inflatus* Kessel 1935: 79, not Lacordaire 1863]. Brazil.
- Entimus* Germar 1817: 341 [= *Curculio* Blanchard 1845: 99, misident., not Linnaeus 1758;
 Entymus Jacob 1937: 155, lapsus]
E. arrogans Pascoe 1872: 448 [= *E. plebejus* Roelofs 1875: XXXVIII]. Colombia, Costa Rica,
 Nicaragua, and Panama.
E. excelsus Viana 1968: 103. Brazil.
E. fastuosus (Olivier 1790: 524) [= *E. splendidus* (Fabricius 1792: 448)]. Brazil.
E. formosus Viana 1958: 2. Argentina.
E. granulatus (Linnaeus 1758: 386) [= *E. sumptuosus* (Olivier 1790: 525); *E. speciosus* Erichson
 1847: 127]. Bolivia, Brazil, Colombia, Ecuador, French Guiana, Guyana, Panama, and
 Peru.
E. imperialis (Forster 1771: 34). Argentina, Brazil, and Paraguay.
E. nobilis (Olivier 1790: 525). Brazil.
E. sastrei Viana 1958: 6 [= *E. nobilis* of authors, misident., not Olivier 1790]. Argentina and
 Uruguay.

Eucoleocerus Champion 1911: 309
E. conicicollis Champion 1911: 309. Mexico.

E. fuscovarius Champion 1911: 310. Guatemala, Honduras, Mexico, and Nicaragua.
E. guttularius Champion 1911: 310. Mexico.

Eudiagogus Schoenherr 1840: 307
E. episcopalis (Gyllenhal 1834: 164). Argentina, Bolivia, Brazil, Ecuador, Paraguay, and Uruguay.
E. pallidevittatus Lucas 1857: 158. Peru.
E. pulcher Fahraeus 1840: 310 [= *E. pogo* Warner 1979: 309]. *Costa Rica, Honduras, and Mexico.

Eudius Schoenherr 1834: 162
E. jocosus Fahraeus 1840: 441. Brazil.
E. quadrisignatus Gyllenhal 1834: 163. Brazil.

Eudmetus Jekel 1854: I.9 bis [= *Eudmetes* Lacordaire 1863: 266, lapsus]
E. cinereus Jekel 1857: 107. Colombia.
E. nigromaculatus Jekel 1857: 111. Colombia.
E. posticatus Jekel 1857: 109. Colombia.

Eurylobus Schoenherr 1826: 152
E. borealis Kessel 1935: 111 [= *E. atromaculatus* Bondar 1949: 178]. Brazil.
E. cinctellus Boheman 1834: 274. Brazil.
E. cingulatus (Germar 1824: 377) [= *E. canephorus* (Germar 1824: 376); *E. fasciatus* Boheman 1834: 274]. Brazil.
E. praecox Kessel 1937: 183. Venezuela (?).
E. surdus Marshall 1949: 331. Brazil.

Euryomus Jekel 1854: I.9
E. affinis Kessel 1935: 106. Brazil.
E. albomaculatus Kessel 1935: 107. Brazil.
E. bondari Marshall 1949: 328. Brazil.
E. chrysiventris Kessel 1935: 103. Brazil.
E. duponti Kessel 1935: 104. Brazil.
E. elegans (Kirby 1819: 437) [= *E. kirbyi* (Gyllenhal 1834: 275)]. Brazil.
E. funebris Kessel 1935: 107. Brazil.
E. griseus Kessel 1935: 105. Brazil.
E. jekeli Kessel 1935: 102. Brazil.
E. mannerheimii (Fahraeus 1842: 185). Brazil and Paraguay.
E. nebulosus Kessel 1935: 108. Brazil.
E. roeselii (Boheman 1842: 184). Brazil.
E. scriptus Kessel 1935: 104. Brazil.
E. ventralis Marshall 1949: 328. Brazil.

Eurysaces Schoenherr 1840: 313
E. grammicus Fahraeus 1840: 314. Brazil.
E. lineolatus (Chevrolat 1881: XXXVIII). Brazil.

Euscapus Jekel 1854: I.6
E. rotundicollis (Boheman 1834: 258). Brazil.
E. squamosus (Fahraeus 1842: 149). Brazil.

Euthyreus Jekel 1854: I.9 bis

E. candidatus (Lacordaire 1863: 269) [= *E. candidatus* Kessel 1935: 78, not Lacordaire 1863].
Brazil.

Eutypus Marshall 1946: 94 [= *Eueides* Jekel 1854: I.7, not Huebner 1816; *Euides* Gemminger
& Harold 1871: 2324, not Fieber 1866 unjust. emend.]

E. chevrolati (Fahraeus 1842: 170). Brazil.

E. clericus (Sahlberg 1823: 34) [= *E. brasiliensis* (Dejean 1821: 91), nom. nud.; *E. scenicus*
(Germar 1824: 373); *E. villosopunctatus* (Kessel 1932: 54), **syn. nov.**]. Brazil.

E. mellyi (Kessel 1932: 52). Brazil.

E. sahlbergi (Fahraeus 1842: 171) [= *E. pupillatus* (Kessel 1932: 51), **syn. nov.**]. Brazil.

Granadia Kessel 1935: 95

G. humeralis Kessel 1935: 96. Colombia.

Hemieuryomus Kessel 1935: 98 [not available, type species not fixed]

H. denudatus Kessel 1935: 100. Brazil.

H. intermedius Kessel 1935: 99. Brazil.

H. magnus Kessel 1935: 99. Brazil.

H. uniformis Kessel 1935: 98. Brazil.

Hypoptophila Voss 1934: 102

H. munda Voss 1934: 102. Colombia.

Hypoptus Jekel 1854: I.9 bis [= *Hypoptoides* Kessel 1932: 59]

H. arcticus Kessel 1932: 62. Colombia.

H. curvipes Voss 1932: 42. Ecuador.

H. gracilis Kessel 1932: 60. Guatemala.

H. immaculatus Kessel 1932: 63. Unknown.

H. insularis Champion 1911: 302. Grenada and St. Vincent.

H. jekeli Champion 1911: 344. Mexico.

H. lepyroides Kessel 1932: 63 [= *H. lepyroides* Lacordaire 1863: 272, nom. nud.]. Colombia.

H. macularis Champion 1911: 303 [= *H. cupreus* Kessel 1932: 62, **syn. nov.**; *H. griseus* Kessel
1932: 62, **syn. nov.**]. Belize, Colombia, Costa Rica, El Salvador, Guatemala, Mexico,
Nicaragua, and Panama.

H. metallicus Kessel 1932: 59. Unknown.

H. punctulatus Kessel 1932: 65. Bolivia.

H. setosulus Kessel 1932: 64. Colombia.

Hypsonotolobus Kessel 1937: 177

H. atlanticus Kessel 1937: 177. Brazil.

Hypsonotus Germar 1824: 367 [= *Charanus* Steven 1829: 92, nom. nud.; *Paraspodus* Jekel
1854: I.9 bis; *Euschoinus* Jekel 1857: 138; *Oligocyrtus* Jekel 1857: 142; *Bactrius* Pascoe
1881: 95; *Curvatonotus* Kessel 1937: 116; *Euscoinus* Kessel 1937: 116, lapsus; *Gibbonotus*
Kessel 1937: 116, not available, type species not fixed; *Punctatonotus* Kessel 1937: 116,
not available, type species not fixed; *Hypsonotus* Vernalha 1952: 34, lapsus]

H. acutipennis Jekel 1859: 229 [= *H. acutipennis* Jekel 1857: 160, nom. nud.]. Colombia.

- H. albicans* Jekel 1857: 156. Brazil.
H. albofasciatus Jekel 1857: 169. Brazil.
H. albopictus Jekel 1857: 181. Brazil.
H. albopunctatus Lucas 1857: 162. Brazil.
H. amazonicus Kessel 1937: 159. Brazil.
H. amoenus Jekel 1857: 155. Brazil.
H. apicalis Kessel 1937: 169. Brazil.
H. apicatus Jekel 1857: 173. Colombia.
H. apocytoides Jekel 1857: 161. Venezuela.
H. aurarius Jekel 1857: 147. Brazil.
H. auripes Jekel 1857: 149 [= *H. auripes* Kessel 1937: 148, not Jekel 1857]. Brazil.
H. auritus Boheman 1834: 257. Brazil.
H. bahianus Kessel 1937: 145. Brazil.
H. binotatus Fahraeus 1842: 163. Brazil.
H. bipunctatus Jekel 1859: 233 [= *H. bipunctatus* Jekel 1857: 160, nom. nud.]. Colombia.
H. bohemanni Jekel 1857: 148. Brazil.
H. boliviensis Kessel 1937: 160. Bolivia.
H. brasiliensis Kessel 1937: 166 [= *H. nigricans* Kessel 1937: 165, **syn. nov.**]. Brazil.
H. brunneus Kessel 1937: 141. Brazil.
H. caliginosus Fahraeus 1842: 155 [= *H. ater* Perroud 1853: 498]. Brazil.
H. callosicollis Jekel 1857: 179 [= *H. includens* Jekel 1857: 183; *H. includens* Lucas 1857: 161, not Jekel 1857]. Brazil and Colombia.
H. camelus Kessel 1937: 127. Brazil.
H. carinatus Kessel 1937: 132. Brazil.
H. cervinus Jekel 1857: 148. Brazil.
H. chrysendetus Germar 1824: 375 [= *H. isabellinemarginatus* Boheman 1834: 260; *H. lateralis* Boheman 1834: 264]. Brazil.
H. cinctipes Boheman 1834: 259 [= *H. reichei* Perroud 1853: 501, **syn. nov.**; *H. arcuatus* Jekel 1859: 203, **syn. nov.**; *H. ater* Jekel 1859: 203, not Perroud 1853; *H. dejeanii* Jekel 1859: 203, nom. nud.]. Brazil.
H. clavulus Germar 1824: 369. Argentina and Brazil.
H. compressipennis Jekel 1859: 235. Colombia.
H. corvinus Kessel 1937: 150. Brazil.
H. dalmani Boheman 1834: 263 [= *H. dalmanni* Jekel 1857: 148, lapsus]. Brazil.
H. decurvatus Kessel 1937: 131. Brazil.
H. dorsalis Fahraeus 1842: 152. Brazil.
H. dorsiger Jekel 1857: 147. Brazil.
H. douei Jekel 1857: 155. Brazil.
H. dromedarius Kessel 1937: 126. Brazil.
H. dubius Kessel 1937: 144. Brazil.
H. fahraei Fahraeus 1842: 154. Brazil.
H. faldermanni Sahlberg 1842: 165. Brazil.
H. fatuus Taschenberg 1907: 226. Brazil.
H. femoralis Kessel 1937: 139. Brazil.
H. frontalis Marshall 1933: 61. Brazil.
H. gibber Kessel 1937: 128. Brazil.
H. glaber Kessel 1937: 170. Colombia.

- H. gounellei* Hustache 1946: 5. Brazil.
H. guatemalenus Kessel 1937: 158. Guatemala.
H. helleri Kessel 1937: 129. Brazil.
H. hondurensis Kessel 1937: 161. Colombia and Honduras.
H. hypocrita Kessel 1937: 165. Brazil.
H. infernalis Kessel 1937: 137. Brazil and Paraguay.
H. interior Kessel 1937: 171. Colombia.
H. interruptilineatus Jekel 1857: 148 [= *H. interruptolineatus* Jekel 1857: 148, lapsus]. Argentina.
H. irrigatus Germar 1824: 375. Brazil.
H. jekelianus Kessel 1937: 156. Brazil.
H. labilis Kessel 1937: 149. Brazil.
H. laevicollis Jekel 1859: 227 [= *H. laevicollis* Jekel 1857: 160, nom. nud.; *H. laevigatus* Jekel 1857: 138, lapsus]. Colombia.
H. languidus Boheman 1859: 123. Ecuador.
H. latissimus Kessel 1937: 125 [= *H. germari* Kessel 1937: 126, **syn. nov.**]. Colombia.
H. latus Jekel 1857: 147. St. Vincent.
H. leucotaenia Kessel 1937: 163. Argentina and Brazil.
H. liber Kessel 1937: 157. Costa Rica.
H. limbifer Kessel 1937: 155. Colombia.
H. lophotoides (Pascoe 1881: 95). Brazil.
H. magnus Kessel 1937: 140. Brazil.
H. melancholicus Jekel 1859: 221 [= *H. melancholicus* Jekel 1857: 159, nom. nud.]. Brazil.
H. mexicanus Kessel 1937: 154. Mexico.
H. mirus Kessel 1937: 167. Brazil.
H. miser Kessel 1937: 135. Colombia or Brazil (?).
H. modestus Marshall 1936: 499. Brazil.
H. molitor Fahraeus 1842: 160 [= *H. incanus* Fahraeus 1842: 157, **syn. nov.**; *H. auriceps* Jekel 1857: 150]. Brazil.
H. morio Jekel 1857: 175. Venezuela.
H. murinus Sahlberg 1842: 162. Venezuela.
H. nebulosus Jekel 1857: 156. Brazil.
H. niger Jekel 1857: 159. Brazil.
H. nigerrimus Kessel 1937: 152 [= *H. lugens* Kessel 1937: 119]. Brazil.
H. nitidulus Jekel 1859: 237 [= *H. nitidulus* Jekel 1857: 160, nom. nud.]. Colombia.
H. nubilosus Fahraeus 1842: 158. Brazil.
H. obsoletus Jekel 1857: 155. Brazil and Colombia.
H. oocularis Jekel 1857: 147. Argentina.
H. ovalis Kessel 1937: 135. Brazil.
H. paraguayanus Jekel 1859: 209 [= *H. paraguayanus* Jekel 1857: 155, nom. nud.]. Paraguay.
H. parceguttatus Marshall 1936: 498. Brazil.
H. parvus Kessel 1937: 167. Brazil.
H. paulistanus Kessel 1937: 164. Brazil.
H. pavidus Jekel 1857: 147. Brazil.
H. phaeopterus Jekel 1857: 149. Brazil.
H. picturatus Sahlberg 1842: 150. Brazil.
H. pilosulus Jekel 1857: 149. Brazil.
H. pulcher Kessel 1937: 143. Brazil.

- H. punctum* Jekel 1859: 231 [= *H. punctum* Jekel 1857: 160, *nom. nud.*]. Colombia.
H. quadrinotatus Kessel 1937: 131. Brazil.
H. ramosus Jekel 1859: 225 [= *H. ramosus* Jekel 1857: 160, *nom. nud.*]. Colombia.
H. rhombifer Marshall 1925: 282. Brazil.
H. rotundipennis Jekel 1857: 167. Brazil.
H. selectus Jekel 1857: 171. Brazil.
H. setarius Jekel 1857: 156. Colombia.
H. setulosus Kessel 1937: 164. Brazil.
H. similis Kessel 1937: 133. Brazil.
H. solutus Kessel 1937: 136. Brazil.
H. stanneus Boheman 1834: 262. Brazil.
H. stevensi Jekel 1857: 165. Brazil.
H. subfasciatus Fahraeus 1842: 159. Brazil.
H. subscutellaris Jekel 1857: 177. Brazil.
H. subvittatus Kessel 1937: 142. Brazil.
H. suturalis Kessel 1937: 138. Brazil.
H. thrichicollis Kessel 1937: 156. Unknown.
H. triangularis Kessel 1937: 134. Brazil.
H. tuberculatus Kessel 1937: 153. Costa Rica.
H. umbilicatus Faust 1892: 28. Venezuela.
H. umbrosus Germar 1824: 370. Brazil.
H. venezolanus Kessel 1937: 159. Venezuela.
H. vestitus Jekel 1859: 223 [= *H. vestitus* Jekel 1857: 159, *nom. nud.*]. Colombia.
H. vilis Kessel 1937: 119. Unknown.
H. viridiadspersus Perroud 1853: 505 [= *H. viridisparus* Jekel 1857: 149]. Brazil.
H. viridipillatus Jekel 1857: 161. Brazil (?).

Lasiocnemoides Wibmer & O'Brien 1986: 93 [= *Lasiocnemus* Jekel 1854: I.7, not Loew 1851]
L. ruficollis (Boheman 1834: 265). Brazil.

L. sculpturatus (Fahraeus 1842: 156) [= *L. niger* (Kessel 1932: 70), not available, color aberration]. Brazil.

Lasiopus Sahlberg 1823: 35 [= *Lasiopus* Schoenherr 1823: col. 1143, *nom. nud.*, *Lasiopus* Schoenherr 1826: 153, not Sahlberg, 1823]

L. cilipes Sahlberg 1823: 35. Brazil.

L. grandis Kessel 1932: 57. Brazil.

L. plumipes (Sahlberg 1842: 172). Brazil.

Lordopella Kessel 1937: 173 [not available, type species not fixed]

L. mirabilis Kessel 1937: 173. Brazil.

L. paraguayana Kessel 1937: 174. Paraguay.

L. parva Kessel 1937: 174. Brazil.

Lordops Schoenherr 1823: col. 1142

L. adspersus Jekel 1854: VI.1. Brazil.

L. affinis Lucas 1857: 164. Brazil.

L. albicinctus Kessel 1932: 9. Brazil.

- L. albofasciatus* Jekel 1854: IV.11. Brazil.
L. alboguttatus Sahlberg 1842: 174. Brazil.
L. amoenus Jekel 1854: IV.5. Brazil.
L. arcuatus Jekel 1854: IV.9. Brazil.
L. argentosus Jekel 1854: IV.15. Brazil.
L. aurosus (Germar 1824: 372). Brazil.
L. brevicornis Kessel 1932: 14. Brazil and Paraguay.
L. catharinensis Jekel 1854: III.11. Brazil.
L. cinctus Fahraeus 1842: 179. Brazil.
L. circumdatus Sahlberg 1842: 176. Brazil.
L. conjugatus Jekel 1854: V.7. Brazil, Colombia, and Paraguay.
L. costifer Jekel 1854: V.13. Brazil.
L. decoratus Kessel 1932: 25. Brazil.
L. deyrollei Jekel 1854: III.9. Brazil.
L. divinus Schenckling & Marshall 1931: 48 [= *L. dives* Lucas 1857: 163, not Germar 1824].
Brazil.
L. elegans Kessel 1932: 8. Brazil.
L. ellipticus Kessel 1932: 7. Brazil.
L. festivus Fahraeus 1842: 180. Brazil.
L. funerarius Jekel 1854: III.15. Brazil.
L. guttatus Fahraeus 1842: 181 [= *L. quinqueguttatus* Jekel 1854: II.7, lapsus]. Brazil.
L. gyllenhalii (Dalman 1823: 83) [= *L. dives* (Germar 1824: 371)]. Brazil.
L. immarginatus Kessel 1932: 15. Brazil.
L. infacetus Boheman 1834: 272. Brazil.
L. insignis Boheman 1834: 269. Brazil.
L. intermedius Kessel 1932: 8. Brazil.
L. interruptus Kessel 1932: 9 [= *L. interuptus* Kessel 1932: 4, lapsus]. Brazil.
L. jekeli Heller 1921: 31 [= *L. transversus* Kessel 1932: 13]. Argentina, Brazil, and Paraguay.
L. jucundus Boheman 1834: 271. Brazil.
L. kuntzeni Kessel 1932: 14. Paraguay.
L. lacrimosus Fahraeus 1842: 175 [= *L. lacrymosus* Jekel 1854: II.4, lapsus]. Brazil.
L. lateralis Fahraeus 1842: 182. Brazil.
L. limbalis Marshall 1949: 165. Brazil.
L. marginatus Jekel 1854: IV.7. Brazil.
L. navicularis (Germar 1824: 371) [= *L. fasciatus* Jekel 1854: V.9, **syn. nov.**]. Brazil.
L. nitidomaculatus Kessel 1932: 17. Brazil.
L. obliquemarginatus Jekel 1854: IV.13. Brazil and Venezuela.
L. obscurus Jekel 1854: V.5. Brazil.
L. ostentatus Jekel 1854: V.3. Brazil.
L. pertyi Jekel 1854: III.7 [= *L. principalis* Perty 1832: 74, misident., not Germar 1824].
Brazil.
L. posticus Jekel 1854: V.1. Brazil.
L. prodigus Jekel 1854: III.5. Brazil.
L. pulvinatus Kessel 1932: 28. Brazil.
L. puncticollis Kessel 1932: 11. Brazil.
L. schoenherri (Dalman 1823: 84) [= *L. principalis* (Germar 1824: 373); *L. buquetii* Jekel 1854:
III.1]. Brazil.

- L. scutellifer* Kessel 1932: 23. Brazil.
L. selectus Kessel 1932: 19. Argentina, Brazil, and Paraguay.
L. semicinctus Jekel 1854: IV.3. Brazil.
L. seminudus Kessel 1932: 8. Brazil.
L. sommeri Fahraeus 1842: 178 [= *L. farinosus* Jekel 1854: V.15; *L. modestus* Jekel 1854: V.16;
 L. V-album Jekel 1854: V.15; *L. variabilis* Jekel 1854: V.15]. Brazil.
L. spinolae Jekel 1854: III.3. Brazil.
L. splendens Kessel 1932: 29. Mexico.
L. transversepunctatus Jekel 1854: III.13. Brazil.
L. tricinctus Jekel 1854: IV.1. Brazil.
L. troberti Jekel 1854: V.11. Brazil.
L. viridulus Boheman 1834: 272. Brazil.
L. waterhousei Jekel 1854: II.15. Brazil.
- Merodontus* Jekel 1854: I.9 bis
M. alophoides Jekel 1857: 129. French Guiana.
M. aurulans Jekel 1857: 123. "Rio Negro".
M. cordifer Jekel 1857: 127. French Guiana.
M. derosus Jekel 1857: 117. French Guiana.
M. forniciatus Jekel 1857: 119. Brazil.
M. setosulus Jekel 1857: 121. Brazil.
M. vetustus Jekel 1857: 125. French Guiana.
- Microstictius* Kuschel 1955: 300
M. naevius (Pascoe 1886: CLI). Brazil.
- Nesolordops* Marshall 1934: 627
N. myersi Marshall 1934: 627. Cuba.
- Omoionotus* Jekel 1854: I.9 bis [= *Homoeonotus* Gemminger & Harold 1871: 2326, unjust.
 emend.]
O. jekeli Faust 1892: 29. Venezuela.
O. lacordairei Faust 1892: 30. Venezuela.
O. miser Kessel 1937: 178. Brazil.
- Orthocnemus* Jekel 1854: I.9 bis
O. heilipoides Jekel 1857: 135. Ecuador and Peru.
O. lebasii Jekel 1857: 133. Colombia.
- Pachyconus* Kessel 1937: 175 [not available, type species not fixed]
P. bahianus Kessel 1937: 176. Brazil.
P. illustris Kessel 1937: 175. Brazil.
- Phaedropus* Schoenherr 1823: col. 1140
P. albobrunneus Bovie 1907: 70. "Guyane".
P. candidus (Fabricius 1775: 146) [= *P. tomentosus* (Olivier 1790: 536); *P. tomentiger* (Herbst
 1795: 319), lapsus]. Brazil and French Guiana.

P. inopinatus Vanin 1983: 295. Bolivia.

P. suturellus Vanin 1983: 297. Brazil.

P. togatus Boheman 1833: 647. Brazil.

Polyteles Schoenherr 1833: 452 [= *Polydius* Schoenherr 1826: 82, not Dejean 1821; *Polytelidius* Bovie, 1908: 43]

P. atrox (Germar 1824: 438). Brazil.

P. decussatus Pascoe 1870: 441. Peru.

P. guerini Fahraeus 1840: 743. Argentina, Bolivia, Brazil, Paraguay, and Uruguay.

P. inka (Heller 1932: 2). Ecuador and Peru.

P. setosus Kirsch 1874: 393. Peru.

P. stevenii (Schoenherr 1826: 82) [= *P. coelestinus* (Perty 1832: 70); *P. orbignyi* Chevrolat 1878: XX]. Argentina, Bolivia, Brazil, and Paraguay.

P. uniformis (Heller 1932: 2). Bolivia.

Pororhynchus Schoenherr 1840: 311 [= *Pororrhynchus* Gemminger & Harold 1871: 2374, lapsus]

P. albolateralis Hustache 1924: 159 [= *P. humeralis* Voss 1934: 102]. Argentina, Bolivia, and Paraguay.

P. labeonis Fahraeus 1840: 312. Argentina, Bolivia, Paraguay, and Uruguay.

P. simplex Voss 1934: 101. Argentina.

Prasocella Kessel 1932: 36

P. peruviana Kessel 1932: 36. Peru.

Promecops Sahlberg 1823: 30 [= *Promecops* Schoenherr 1823: col. 1141; *Periorges* Schoenherr 1845: 420; *Promecopsis* Champion 1911: 312, not Duméril 1806; *Exeudius* Voss 1934, not available, type species not fixed; *Promecodes* Marshall 1939: 582, not available, type species not fixed; *Promecodes* Balfour-Browne 1960: 240]

P. aemulus Voss 1934: 84. Bolivia and Peru.

P. affinis Hustache 1924: 157. Bolivia.

P. alternans Champion 1911: 316. Belize, Costa Rica, Guatemala, Honduras, and Mexico.

P. angustifrons Voss 1934: 87. French Guiana.

P. arcuatus Fahraeus 1840: 329 [= *P. nitidiventris* (Lucas 1857: 160)]. Brazil.

P. bahianus Voss 1939: 339. Brazil.

P. boliviarius Voss 1934: 90 [= *P. similaris* Voss 1934: 86]. Bolivia and Brazil.

P. boops Gyllenhal 1834: 165. Brazil.

P. brevisetis Champion 1911: 316. Guatemala and Mexico.

P. campanulicollis Voss 1934: 88. Bolivia and Brazil.

P. carinellus Hustache 1923: 299. Bolivia.

P. carinicollis Hustache 1923: 296 [= *P. brevis* Hustache 1923: 297; *P. amazonica* Voss 1934: 89]. Bolivia, Brazil, and French Guiana.

P. carinirostris Voss 1934: 82. Costa Rica.

P. cerussatus Fahraeus 1840: 319 [= *P. semicarinulata* Voss 1934: 85; *P. subsetosa* Voss 1934: 87; *P. ingae* Voss 1951: 369]. Brazil.

P. claviger Hustache 1924: 156. Bolivia.

P. clavisetis Champion 1911: 315. Honduras and Mexico.

- P. clitellarius* Fahraeus 1840: 334. Brazil and Paraguay.
- P. concinnus* Voss 1934: 76. Peru.
- P. consobrinus* Voss 1953: 433. Brazil.
- P. costaricensis* Voss 1934: 80. Costa Rica.
- P. curvirostris* Voss 1934: 77 [= *P. sieberi* Voss 1934: 77]. Bolivia, Brazil, and Peru.
- P. dentimanus* Champion 1911: 314. Panama.
- P. diversus* Voss 1934: 96. Brazil.
- P. francoi* Voss 1953: 431 [= *P. novaeutoniae* Voss 1953: 428]. Brazil.
- P. gracilis* Fahraeus 1840: 336 [= *P. pulchellus* Boheman 1859: 122]. Brazil.
- P. impuratus* Gyllenhal 1834: 167. Brazil.
- P. irroratus* (Lucas 1857: 161). Brazil.
- P. jucundus* Faust 1892: 26. Venezuela.
- P. lepidoides* Voss 1934: 75. Colombia.
- P. lepidus* Fahraeus 1840: 317. Colombia.
- P. leucothyreus* Fahraeus 1840: 328. Bolivia, Colombia, Costa Rica, El Salvador, Guatemala, Honduras, Mexico, and Nicaragua.
- P. limbatus* Gyllenhal 1834: 168 [= *P. dorsalis* Fahraeus 1840: 335; *P. fallax* Fahraeus 1840: 333; *P. peruana* Voss 1934: 96]. Argentina, Brazil, and Peru.
- P. lineiger* Pascoe 1886: CLV. Brazil.
- P. lunatus* Fahraeus 1840: 327 [= *P. cognatus* Fahraeus 1840: 331]. Grenadines and St. Vincent.
- P. luteus* Faust 1892: 25 [= *P. laetus* Faust 1892: 26; *P. luneus* Blackwelder 1947: 806, lapsus]. Brazil.
- P. modestus* Voss 1934: 94. Brazil.
- P. nebulosus* Kirsch 1874: 394 [= *P. infidus* Kirsch 1874: 395; *P. solidus* Voss 1934: 76]. Peru.
- P. nubeculosus* Gyllenhal 1834: 166 [= *P. cinctus* Fahraeus 1840: 323; *P. petulans* Voss 1934: 89; *P. canavaliae* (Marshall 1937: 471)]. Brazil and French Guiana.
- P. nubifer* Sahlberg 1823: 30. Brazil.
- P. obscurus* Voss 1934: 93. Brazil and French Guiana.
- P. ohausi* Voss 1934: 79. Ecuador.
- P. parvulus* Fahraeus 1840: 332. Brazil.
- P. pauxillus* Voss 1934: 78. Brazil.
- P. phaleratus* Lucas 1857: 158. Brazil.
- P. posticus* Fahraeus 1840: 322 [= *P. alboscutellatus* (Chevrolat 1880: 254)]. Guadeloupe.
- P. proximus* Voss 1934: 89. Brazil.
- P. puncticollis* Hustache 1923: 298 [= *P. irroratus* Voss 1934: 95, not Lucas 1857; *P. leucostictus* Voss 1934: 95]. Argentina, Bolivia, Brazil, and Paraguay.
- P. rhombicus* Fahraeus 1840: 326. Colombia.
- P. rhombifera* Fahraeus 1840: 325 [= *P. olivieri* Faust 1892: 27; *P. stimulans* Faust 1892: 28]. Colombia, Guyana, and Venezuela.
- P. setiger* Fahraeus 1840: 318 [= *P. umbrosus* Voss 1934: 81]. Brazil.
- P. setosus* Voss 1934: 86. Brazil.
- P. sinuatocollis* Hustache 1929: 202. Guadeloupe.
- P. sitonoides* Voss 1953: 428. Brazil.
- P. spathula* Voss 1939: 338. Brazil.
- P. subsignatus* (Boheman 1845: 421) [= *P. rostralis* Voss 1934: 82]. Brazil.
- P. torvus* Fahraeus 1840: 330. Brazil.
- P. tridentatus* Champion 1911: 315. Guatemala.

- P. umbratus* Fahraeus 1840: 324 [= *P. viator* Fahraeus 1840: 332]. Mexico.
P. unidentatus Champion 1911: 313. Costa Rica (?) and Panama.
P. uniformis Champion 1911: 317. Guatemala.
P. vicinus Hustache 1924: 158. Bolivia.
P. vulpinus Fahraeus 1840: 316 [= *P. scrobicollis* Fahraeus 1840: 320; *P. vagabundus* Fahraeus 1840: 321]. Brazil.
- Pseudhypoptus* Champion 1911: 304
P. eurylobus Champion 1911: 305. Mexico.
P. parcus (Fahraeus 1842: 183). Mexico.
- Rescapus* Kessel 1935: 74
R. sapiens Kessel 1935: 74. Brazil.
- Rhigus* Schoenherr 1823: col. 1138 [= *Rhigus* Germar 1824: 438, not Schoenherr 1823]
R. agricola Boheman 1840: 732. Brazil.
R. dejeanii Gyllenhal 1833: 446 [= *R. contaminatus* Gyllenhal 1833: 446]. Brazil.
R. faldermanni Boheman 1840: 733 [= *R. irroratus* Boheman 1840: 735, **syn. nov.**]. Argentina, Brazil, and Paraguay.
R. horridus Dalman 1833: 445. Brazil.
R. lateritus Gyllenhal 1833: 445. Brazil.
R. nigrosparsus (Perty 1832: 71) [= *R. multipunctatus* Boheman 1840: 734]. Argentina, Brazil, and Paraguay.
R. phaleratus Perty 1832: 71 [= *R. gyllenhalii* Boheman 1833: 448]. Brazil.
R. schueppelii Germar 1824: 439. Brazil.
R. speciosus (Linnaeus 1758: 385) [= *R. smaragdulus* (Fabricius 1787: 111); *R. bidentatus* Boheman, 1833: 448]. French Guiana.
R. tribuloides (Pallas 1781: 30) [= *R. brunneus* Bovie 1907: 328; *R. bunneus* Bovie 1908: 5, lapsus]. Brazil and Paraguay.
R. vespertilio Pascoe 1881: 299 [= *R. tribuloides* Bovie 1908: Fig. 13, misident., not Pallas 1781]. Argentina and Brazil.
- Stenorhinus* Jekel 1854: I.7 [= *Stenorhinus* Schenkling & Marshall 1931: 54, lapsus]
S. leucostictus (Germar 1824: 374). Brazil.
S. viridimarginatus (Boheman 1834: 261). Brazil.
- Sullana* Blackwelder 1947: 808 [= *Sulla* Kessel 1937: 182, not Lesson 1837]
S. columbiana (Kessel 1937: 118). Colombia.
- Tomometopus* Jekel 1854: I.9 bis
T. lordopiooides Kessel 1932: 67 [= *T. lordopiooides* Lacordaire 1863: 269, nom. nud., *T. lordopiooides* Gemminger & Harold 1871: 2322, nom. nud.]. Bolivia.
- Tomorhinus* Jekel 1854: I.8
T. designatus (Fahraeus 1842: 164) [= *T. designatus* Kessel 1935: 76]. Brazil.
T. pictus Kessel 1935: 164) [= *T. pictus* Lacordaire 1863: 270, nom. nud.; *T. pictus* Gemminger & Harold 1871: 2323, nom. nud.]. French Guiana.
- Trachyus* Kuschel 1955: 301

T. fischeri (Gyllenhal 1833: 451). Brazil.

Trichocnemus Jekel 1854: VI.3

T. suturalis Jekel 1854: VI.5. Brazil.

Tropidorrhinus Gemminger & Harold 1871: 2325 [= *Tropirhinus* Schoenherr 1840: 360, not Schoenherr 1823; *Tropirrhinus* Schenkling & Marshall 1931: 56, lapsus]

T. affaber (Fahraeus 1840: 361). Brazil.

T. costatus (Boheman 1834: 262) [= *T. costipennis* (Sahlberg 1842: 166); *T. interruptus* (Kessel 1935: 83), **syn. nov.**]. Brazil.

Subtribe OPHRYASTINA Lacordaire 1863

Ophryastes Say 1831: 13 [= *Ophryastes* Schoenherr 1833: 508, not Say 1831; *Dystirus* Pascoe 1872: 447; *Eupagoderes* Horn 1876: 32; *Caccophryastes* Sharp 1891: 92; *Tosastes* Sharp 1891: 91; *Amydrogmus* Pierce 1913: 374]

O. argentatus LeConte 1853: 444. *Mexico.

O. aridus (Fall 1910: 192). *Mexico.

O. cinereus Fahraeus 1840: 819. *Mexico.

O. constrictus (Sharp 1891: 93). Mexico.

O. decipiens LeConte 1853: 445. *Mexico.

O. depressirostris (Champion 1911: 321). Mexico.

O. desertus (Horn 1876: 34) [= *O. giganteus* (Chittenden 1926: 169); *O. californicus* (Ting 1939: 81)]. *Mexico.

O. durangoensis (Sharp 1891: 94). Mexico.

O. globipennis (Sharp 1891: 91). Mexico.

O. gracilis (Sharp 1891: 96). Mexico.

O. huachucae (Van Dyke 1934: 180). *Mexico.

O. latirostris LeConte 1853: 443 [= *O. validus* LeConte 1856 225; *O. strumosus* (Pascoe 1872: 447); *O. sulcipennis* Casey 1888: 239; *O. latipennis* Sharp 1891: 89]. *Mexico.

O. lineatus (Sharp 1891: 92). Mexico.

O. lucanus (Horn 1876: 34). Mexico.

O. mexicanus (Sharp 1891: 94). Mexico.

O. neogracilis Kissinger 1970: 47. Mexico.

O. ovipennis Sharp 1891: 90 [= *O. bituberosus* Pierce 1909: 344, misident., not Sharp 1891]. *Mexico.

O. porosus LeConte 1854: 225 [= *O. humeralis* (Sharp 1891: 91); *O. coarctatus* (Champion, 1911: 319)]. Mexico.

O. prolatus (Sharp 1891: 94). Mexico.

O. robustus (Davis 1947: 503). *Mexico.

O. rotundatus (Champion 1911: 320). *Mexico.

O. sallei Gyllenhal 1840: 820. Mexico.

O. simulans (Van Dyke 1934: 176). *Mexico.

O. sordidus LeConte 1853: 445 [= *O. cretaceus* (Sharp 1891: 96); *O. halli* (Van Dyke 1934: 181)]. *Mexico.

O. speciosus LeConte 1853: 444. *Mexico.

O. squalidus (Sharp 1891: 95). Mexico.

- O. sulcirostris* (Say 1824: 316) [= *O. ligatus* LeConte 1853: 443; *O. porosus* Davis 1947: 531, misident., not LeConte 1856]. *Mexico.
O. symmetricus Fall 1907: 260. *Mexico.
O. tuberosus LeConte 1853: 443 [= *O. basalis* Sharp 1891: 89; *O. bituberosus* Sharp 1891: 90; *O. collaris* Champion 1911: 319; *O. tetralobus* Champion 1911: 318]. *Mexico.
O. turbinatus (Champion 1911: 320). *Mexico.
O. variabilis (Pierce 1913: 374). *Mexico.
O. varius LeConte 1853: 444. *Mexico.
O. vittatus (Say 1824: 316). *Mexico.

Subtribe EUSTYLINA Lacordaire 1863

Exophthalmina Champion 1911.
Exophthalmodina Voss 1954.

- Anidius* Kuschel 1955: 305
A. duricornis (Gyllenhal 1840: 367). Brazil.
A. fallax (Gyllenhal 1840: 368). Brazil.
A. lepidotus (Bohemian 1840: 366). South America.
A. duricornis (Gyllenhal 1840: 367). Brazil.
A. nitidulus (Fabricius 1775: 146) [= *A. circumductus* (Germar 1824: 454); *A. auronitens* (Mannerheim 1833: 650); *A. viridisquamulosus* (Bohemian 1840: 376)]. Brazil and French Guiana.

- Aphrastus* Say 1821: 9 [= *Aphrastus* Schoenherr 1833: 16, not Say 1831; *Micronychus* Provancher 1877: 508]
A. angularis Champion 1911: 299. Mexico.
A. submarginatus Champion 1911: 299. Mexico.

- Brachyomus* Lacordaire 1863: 130 [= *Cyphipterus* Dejean 1835: 271, 1837: 294, nom. nud.]
B. bicostatus Faust 1892: 14. Venezuela.
B. histrio Faust 1892: 14. Venezuela.
B. metallescens Pascoe 1880: 427. Ecuador and Peru.
B. octotuberculatus (Fabricius 1787: 112). French Guiana, Trinidad, and Venezuela.
B. quadrinodosus (Bohemian 1842: 217) [= *B. sallei* Faust 1892: 15]. Colombia and Venezuela.
B. quadrifibulatus (Bohemian 1842: 216). Colombia and Venezuela.
B. tuberculatus (Bohemian 1842: 218). St. Vincent.
Brachystylus Schoenherr 1845: 433 [= *Syntomostylus* Scudder 1893: 50, unjust. replacement name]
B. microphthalmus Champion 1911: 298. Belize and Mexico.
B. tamaulipanus Champion 1911: 298. Mexico.

- Compsus* Schoenherr 1823: col. 1140 [= *Compsus* Schoenherr 1826: 109, not Schoenherr 1823; *Comlabus* Voss 1953: 57, lapsus; *Percompsus* Voss 1954: 203]
C. acrolithus (Germar 1824: 435). Brazil.
C. adonis Marshall 1922: 197. Colombia.
C. aeruginosus (Bohemian 1840: 177). Colombia.
C. affinis Hustache 1938: 77. Colombia.

- C. ainaensis* Voss 1954: 203. Peru.
- C. albosetosus* Hustache 1938: 84 [= *C. albosetotus* Hustache 1938: 71, lapsus]. Colombia.
- C. albus* Hustache 1938: 76. Colombia.
- C. alternans* Kirsch 1889: 20. Ecuador.
- C. alternevittatus* Hustache 1938: 83. Colombia.
- C. argyreus* (Linnaeus 1758: 384) [= *C. elegans* (Olivier 1807: 328); *C. litargyreus* (Fauvel 1861: 314), lapsus; *C. subelongatus* Voss 1954: 202, **syn. nov.**]. Argentina, Bolivia, Brazil, Ecuador, French Guiana, Peru, and Suriname.
- C. armatissimus* Marshall 1922: 193 [= *C. serietuberculatus* Voss 1932: 38]. Brazil.
- C. ater* Kirsch 1889: 20. Colombia.
- C. attenuatus* Hustache 1938: 84 [= *C. interruptus* Hustache 1938: 84, **syn. nov.**]. Colombia.
- C. auricephalus* (Say 1824: 310) [= *C. auriceps* (Schoenherr 1840: 183)]. Costa Rica, Guatemala, Honduras, Mexico, and Nicaragua.
- C. aurisquamatus* Hustache 1938: 99. Venezuela.
- C. azureipes* Hustache 1938: 87. Brazil.
- C. benoisti* Hustache 1938: 106. Ecuador.
- C. bicarinatus* Kirsch 1889: 22. Colombia.
- C. bicoloratus* (Boheman 1840: 196). Peru.
- C. biumpresus* Faust 1892: 9. Venezuela.
- C. bisignatus* (Boheman 1840: 190). Argentina and French Guiana.
- C. bituberculatus* Kirsch 1889: 19. Colombia.
- C. bituberous* Kirsch 1868: 237. Colombia.
- C. bourcieri* (Guérin 1855: 591). South America.
- C. caeruleipes* Champion 1911: 288. Costa Rica and Panama.
- C. candidus* Hustache 1938: 75. Colombia.
- C. canescens* (Boheman 1840: 181) [= *C. dorsalis* Hustache 1938: 84, **syn. nov.**]. Colombia.
- C. chlorostictus* (Erichson 1848: 568). Guyana.
- C. chrysis* (Fabricius 1792: 449). Neotropical.
- C. clarus* (Fabricius 1801: 541). Guyana.
- C. cometes* (Pascoe 1880: 425) [= *C. adamantinus* Kirsch 1889: 19]. Ecuador.
- C. confluens* Marshall 1922: 194. Brazil.
- C. coruscus* (Pascoe 1880: 425). Ecuador.
- C. costaricensis* Voss 1932: 39. Costa Rica.
- C. cyanipes* (Fabricius 1787: 113). French Guiana.
- C. cyanitarsis* Hustache 1938: 93. Bolivia and Colombia.
- C. cyphoides* Hustache 1938: 108. Colombia.
- C. dalmani* (Boheman 1840: 175) [= *C. dives* (Guérin 1833: pl. 27), misident., not Olivier 1807; *C. dalmanni* (Guérin 1844: 141), lapsus]. Brazil.
- C. delicatulus* Hustache 1938: 79. Colombia.
- C. deliciosus* Hustache 1938: 80. Colombia.
- C. dentipes* Marshall 1949: 326. Brazil.
- C. deplanatus* Kirsch 1868: 237. Colombia and Venezuela.
- C. dives* (Olivier 1807: 296). South America.
- C. divisus* Hustache 1938: 82. Colombia.
- C. ermineus* (Boheman 1840: 191) [= *C. molitor* (Rosenschoeld 1840: 189); *C. atrigutta* Marshall 1949: 325]. Argentina, Bolivia, Brazil, and Paraguay.
- C. eustyloides* Hustache 1938: 96 [= *C. conspicillatus* Hustache 1938: 98, **syn. nov.**]. Colombia.

- C. exanguis* (Bohemian 1833: 641) [= *C. exsanguis* (Schoenherr 1840: 202), lapsus; *C. virgineus* Pascoe 1880: 423]. Peru.
- C. fossicollis* Kirsch 1889: 19. Ecuador.
- C. fractilineatus* Marshall 1922: 200. Venezuela.
- C. fulgidipes* Bovie 1907: 326. Brazil.
- C. furcatus* (Guérin 1855: 592) [= *C. fuscatus* Lacordaire 1863: 117, lapsus]. South America.
- C. gemmeus* Faust 1892: 13. Venezuela.
- C. gentilis* (Olivier 1807: 315). Guadeloupe.
- C. glaucus* (Bohemian 1840: 178). Colombia.
- C. iris* Marshall 1922: 198. Colombia.
- C. isabellinus* (Bohemian 1833: 644) [= *C. albarius* (Bohemian 1840: 192)]. Argentina, Bolivia, and Brazil.
- C. lacteus* (Fabricius 1781: 185) [= *C. religiosus* (Olivier 1807: 302), **syn. nov.**; *C. canus* (Schoenherr 1826: 110), **syn. nov.**; *C. dejeani* (Bohemian 1840: 185), **syn. nov.**]. French Guiana, Guadeloupe, and Jamaica.
- C. latifrons* Hustache 1938: 90 [= *C. hypocrita* Hustache 1938: 91, **syn. nov.**; *C. hypocritus* Hustache 1938: 91, lapsus]. Colombia.
- C. lebasii* (Bohemian 1840: 182). Colombia.
- C. lepidus* (Bohemian 1840: 193). French Guiana.
- C. lineatus* Hustache 1938: 81. Colombia.
- C. luquillo* Wolcott 1951: 390. Puerto Rico.
- C. maricao* Wolcott 1924: 125. Puerto Rico.
- C. monachus* (Fauvel 1861: 313). French Guiana.
- C. nigropunctatus* Champion 1911: 289. Costa Rica, Guatemala, and Panama.
- C. niveus* (Fabricius 1787: 111) [= *C. otti* Hustache 1938: 74]. Brazil and French Guiana.
- C. obliquatus* Hustache 1938: 77. Colombia.
- C. ochroleucus* (Erichson 1848: 568). Guyana.
- C. opulentus* Kirsch 1889: 18. Ecuador.
- C. ostracion* (Pascoe 1880: 420). Brazil.
- C. parviscutum* Hustache 1938: 94. Ecuador.
- C. parvus* Hustache 1938: 103. Venezuela.
- C. pertinax* Hustache 1938: 85 [= *C. dorsofuscus* Hustache 1938: 86, **syn. nov.**; *C. vilosus* Hustache 1938: 86, **syn. nov.**]. Colombia.
- C. peruvianus* Hustache 1938: 86. Peru.
- C. placidus* (Bohemian 1840: 180). Colombia.
- C. popayanus* Kirsch 1889: 21. Colombia.
- C. pugionatus* Marshall 1922: 196. Colombia, Venezuela, and Central America.
- C. pustulosus* Hustache 1938: 92. Ecuador.
- C. quadrilineatus* Kirsch 1889: 18. Ecuador.
- C. quadrifasciatus* (Bohemian 1840: 188) [= *C. simoni* Faust 1892: 10; *C. octodecimsignatus* Champion 1911: 288; *C. viridipunctatus* Hustache 1938: 78]. Ecuador.
- C. roseomicans* Hustache 1938: 100. Bolivia and Colombia.
- C. rufipes* (Bohemian 1840: 199). Brazil.
- C. saucius* (Bohemian 1840: 198). Brazil.
- C. scrutator* Hustache 1938: 98. Brazil, Colombia, and Venezuela.
- C. sejunctus* (Pascoe 1880: 426) [= *C. interruptus* (Pascoe 1880: 426); *C. variegatus* Kirsch 1889: 21]. Brazil and Ecuador.

- C. serrans* Kuschel 1955: 2. Venezuela.
C. servus (Fabricius 1787: 112) [= *C. alboguttatus* Hustache 1938: 102]. French Guiana.
C. sordidus (Boheman 1840: 174) [= *C. obliquecostatus* (Perroud 1853: 464)]. Brazil.
C. spectabilis Faust 1892: 12. Venezuela.
C. subcostatus Hustache 1938: 100. Venezuela.
C. sulcicollis Hustache 1938: 101. Venezuela.
C. sylvaticus Hustache 1938: 91. Venezuela.
C. tripunctatus Hustache 1938: 105. Ecuador.
C. vespertinus Pascoe 1880: 424. Peru.
C. violaceus Hustache 1938: 89. Colombia.
C. virginalis Faust 1892: 9. Venezuela.
C. viridimaculatus (Boheman 1840: 185). French Guiana.
C. viridis Voss 1932: 38. Costa Rica.
C. viridissimus Hustache 1938: 107. Colombia and Ecuador.
C. viridivittatus (Guérin 1855: 592). Colombia.
C. viridulus Hustache 1938: 95. Colombia.
C. westermanni (Boheman 1840: 197). Brazil.
C. zebra Marshall 1922: 199. Colombia and Ecuador.
C. zebrinus Voss 1953: 57. Colombia.

Diaprepes Schoenherr 1823: col. 1140

- D. abbreviatus* (Linnaeus 1758: 386) [= *D. brevis* (Olivier 1790: 549), **syn. nov.**; *D. festivus* (Fabricius 1792: 476), not Fabricius 1792: 404; *D. irregularis* (Panzer 1798: 67); *D. japonensis* (Voet 1806: 52); *D. festivus* (Olivier 1807: 312), not Fabricius 1792, **syn. nov.**; *D. quadrilineatus* (Olivier 1807: 310), **syn. nov.**; *D. distinguendus* Gyllenhal 1834: 10, **syn. nov.**; *D. vicinus* Dejean 1834: 252, *nom. nud.*; *D. guadeloupensis* Gyllenhal 1834: 11, **syn. nov.**; *D. guadeloupensis* Hustache 1929: 184, **syn. nov.**]. Barbados, Dominican Republic, Guadeloupe, Hispaniola, Martinique, Mona, Montserrat, Puerto Rico, St. Lucia, St. Vincent, and Vieques.
D. albofasciatus Champion 1911: 180. Honduras.
D. balloui Marshall 1916: 449. Dominican Republic.
D. barbadensis Marshall 1916: 451, **stat. nov.** (from subspecies of *D. famelicus*). Barbados.
D. boxi Marshall 1938: 3. St. Lucia.
D. comma Boheman 1834: 8. Trinidad and Venezuela.
D. doublieri Guérin 1846: 428 [= *D. comma* Pierce 1915: 258, misident., not Boheman 1834]. Haiti.
D. elegantulus Gyllenhal 1834: 13, **stat. nov.** (from subspecies of *D. famelicus*). Cuba and Martinique.
D. esuriens Gyllenhal 1834: 15 [= *D. purvesi* Roelofs 1875: XXVI], **stat. nov.** (from subspecies of *D. famelicus*). Antigua, Montserrat, St. Barthélemy, and St. Kitts.
D. excavatus Rosenschoeld 1840: 343. St. Vincent.
D. famelicus (Olivier 1790: 544) [= *D. affinis* (Fabricius 1801: 531); *D. lepidopterus* Gyllenhal 1834: 14; *D. leucopterus* (Leng and Mutchler 1914: 469)]. Dominican Republic, Guadeloupe, Martinique, and Nevis.
D. glaucus (Olivier 1807: 315). West Indies.
D. marginatus (Fabricius 1775: 145) [= *D. bivittatus* (Fabricius 1787: 118); *D. circumdatus* Schoenherr 1826: 117, *nom. nud.*; *D. denudatus* Pierce 1915: 263]. Guadeloupe and St. Thomas.

- D. maugei* (Boheman 1840: 369) [= *D. capsicalis* Marshall 1922: 59]. Puerto Rico.
- D. pulverulentus* Champion 1911: 180. Nicaragua.
- D. reticulatus* Chevrolat 1880: 165. Martinique.
- D. revestitus* Chevrolat 1880: 165. Cuba.
- D. rohrii* (Fabricius 1775: 151) [= *D. lundii* (Herbst 1795: 134); *D. farinosus* Gyllenhal 1840: 344; *D. marginatus* Pierce 1915: 258, misident., not Fabricius 1775]. St. Croix.
- D. rufescens* Boheman 1840: 346. Guadeloupe.
- D. sommeri* (Rosenschoeld 1840: 339). Cuba.
- D. spengleri* (Linnaeus 1767: 609) [= *D. sprengleri* (Linnaeus 1767: 609), incorrect original spelling; *D. sommeri* Guenther and Zumpt 1933: 109, lapsus, in part]. Puerto Rico.
- D. variegatus* Chevrolat 1880: 165. Martinique.
- Eustylus* Schoenherr 1843: 40
- E. aequus* Marshall 1916: 464. Brazil.
- E. bodkini* Marshall 1916: 456. Colombia, Guyana, and Venezuela.
- E. chiriquensis* Champion 1911: 294. Panama.
- E. cinericius* Champion 1911: 296. Guatemala.
- E. ephippiatus* Marshall 1916: 458. Colombia.
- E. funicularis* Kirsch 1874: 390 [= *E. bolivianus* Marshall 1916: 463]. Bolivia and Peru.
- E. grypsatus* (Boheman 1843: 27). Mexico.
- E. hybridus* (Rosenschoeld 1840: 200). Guadeloupe and St. Lucia.
- E. inclusus* Marshall 1916: 461. Brazil.
- E. magdalena* Marshall 1926: 536 [= *E. magdalensis* Lona 1938: 446, lapsus]. Colombia.
- E. obliquefasciatus* Marshall 1916: 455. Venezuela.
- E. puber* (Olivier 1807: 367) [= *E. aparius* (Rosenschoeld 1840: 184)]. Colombia, French Guiana, Guyana, Trinidad, and Venezuela.
- E. quadrigibbus* Champion 1911: 292. Guatemala.
- E. ruptus* Champion 1911: 293. Costa Rica and Panama.
- E. scapularis* Marshall 1916: 464. Brazil.
- E. setipennis* Champion 1911: 292. Guatemala.
- E. sexguttatus* Champion 1911: 291. Panama.
- E. simplex* Marshall 1916: 460. Colombia.
- E. simulatus* Marshall 1916: 459. Colombia.
- E. sordidus* Marshall 1916: 460. Colombia.
- E. striatus* Boheman 1843: 42. Mexico.
- E. subapterus* Champion 1911: 295. Guatemala and Mexico.
- E. subfasciatus* Voss 1932: 40. Paraguay.
- E. subguttatus* Marshall 1916: 462. Brazil.
- E. subvittatus* Marshall 1916: 457. Venezuela.
- E. veraepacis* Champion 1911: 294. Guatemala.
- Exophthalmus* Schoenherr 1823: col. 1140 [= *Prepodes* Schoenherr 1823: col. 1140; *Praepodes* Schoenherr 1840: 348, lapsus; *Exophthalmodes* Pierce 1916: 464]
- E. agrestis* (Boheman 1834: 291) [= *E. canus* Gyllenhal 1834: 306; *E. canescens* (Boheman 1840: 357); *E. farinolentus* (Boheman 1840: 358); *E. uniformis* (Boheman 1842: 213)]. Mexico and Paraguay.
- E. albidus* Champion 1911: 255. Mexico.

- E. albolineatus* Champion 1911: 262. Mexico.
- E. albosquamatus* (Sallé 1855: 268). Dominican Republic.
- E. albovittatus* Champion 1911: 263. Guatemala.
- E. annulonotatus* (Waterhouse 1879: 423). Colombia and Ecuador.
- E. aurarius* (Gyllenhal 1834: 12). Guadeloupe.
- E. bilineatus* Champion 1911: 265. Guatemala.
- E. caeruleovittatus* Champion 1911: 264. Guatemala.
- E. carinirostris* (Bohemian 1840: 187). Guatemala, Mexico, and Nicaragua.
- E. carneipes* Champion 1911: 257. Costa Rica, Honduras, and Panama.
- E. chrysopus* Chevrolat 1880: 165. Dominican Republic and Haiti.
- E. cinerascens* (Fabricius 1792: 452). French Guiana and Hispaniola.
- E. clathratus* Champion 1911: 264. Mexico.
- E. consobrinus* (Marshall 1922: 190). Colombia.
- E. costatus* (Gyllenhal 1834: 18). Hispaniola.
- E. crassicornis* Kirsch 1868: 238. Colombia and Panama.
- E. cupreipes* Champion 1911: 257. Mexico.
- E. cuprirostris* Champion 1925: 427. Costa Rica.
- E. depressicollis* (Gyllenhal 1834: 22). Hispaniola.
- E. distigma* Champion 1911: 259. Guatemala.
- E. dufaui* (Hustache 1929: 198). Guadeloupe.
- E. duplicatus* Champion 1911: 263. Mexico.
- E. farr* Vaurie 1961: 35. Jamaica.
- E. fasciatus* Champion 1911: 252. Mexico.
- E. foveicollis* (Chevrolat 1880: 175). Guadeloupe.
- E. frenatus* (Marshall 1934: 624). Dominican Republic and Haiti.
- E. gundlachi* Chevrolat 1880: 165. Cuba.
- E. haitiensis* Bovie 1907: 68. Haiti.
- E. hemigrammus* (Chevrolat 1880: 197) [= *E. hemipterus* Lona 1938: 520, lapsus]. Martinique.
- E. hieroglyphicus* Chevrolat 1879: XCVII. Hispaniola.
- E. humeridens* (Marshall 1934: 623). Cuba.
- E. humilis* (Gyllenhal 1834: 28). Brazil.
- E. hybridus* (Chevrolat 1858: 210). Cuba.
- E. impositus* (Pascoe 1880: 427). Costa Rica and Nicaragua.
- E. impressus* (Fabricius 1781: 192) [= *E. albescens* (Cockerell 1893: 374); *E. roseomaculatus* (Cockerell 1893: 374)]. Jamaica.
- E. inaequalis* (Gyllenhal 1834: 23). Hispaniola.
- E. interpositus* Champion 1911: 253. Mexico.
- E. interruptus* (Chevrolat 1880: 165). Guadeloupe.
- E. jekelianus* (White 1858: 357) [= *E. stellaris* (Pascoe 1880: 425)]. Colombia, Costa Rica, Nicaragua, and Panama.
- E. laetus* (Olivier 1807: 326). Dominican Republic and Haiti.
- E. lepidus* (Chevrolat 1858: 210). Cuba.
- E. leucographus* (Gyllenhal 1834: 19). Hispaniola.
- E. lunaris* Champion 1911: 266. Belize, Guatemala, and Mexico.
- E. maculosus* Chevrolat 1879: XCVIII. Cuba.
- E. mannerheimi* Boheman 1840: 338. Dominican Republic and Haiti.
- E. margaritaceus* Champion 1911: 255. Honduras and Nicaragua.

- E. marginicollis* (Chevrolat 1880: 175). Guadeloupe.
- E. marmoreus* (Gyllenhal 1840: 352) [= *E. marmoratus* (Hustache 1929: 198), **syn. nov.**].
Guadeloupe.
- E. martinicensis* Chevrolat 1879: XC VIII. Martinique.
- E. mixtus* Chevrolat 1880: 165. Cuba.
- E. murinus* (Rosenschoeld 1840: 354). Suriname.
- E. nicaraguensis* Bovie 1907: 327 [= *E. nicaraguaensis* Lona 1938: 521, lapsus]. Costa Rica,
Nicaragua, and Panama.
- E. nubilus* (Rosenschoeld 1840: 355). Mexico.
- E. obsoletus* (Olivier 1807: 325) [= *E. sejugatus* (Schoenherr 1826: 118)]. Haiti.
- E. opulentus* (Boheman 1840: 365). Guatemala and Mexico.
- E. ornatus* Champion 1911: 253. Guatemala.
- E. parentheticus* (Marshall 1922: 191). Ecuador.
- E. pictus* (Guérin 1847: 9) [= *E. fulvovirgatus* (Marshall 1934: 625), **syn. nov.**]. Cuba.
- E. plicatus* Champion 1911: 268. Panama.
- E. pugnax* (Olivier 1807: 349) [= *E. forstroemi* Gyllenhal 1834: 6]. Haiti.
- E. pulchellus* Boheman 1834: 5. South America.
- E. quadriplagiatus* (Olivier 1807: 325). Dominican Republic and Haiti.
- E. quindecimpunctatus* (Olivier 1807: 300). Puerto Rico.
- E. roralis* (Boheman 1840: 356). Mexico.
- E. roseipes* (Chevrolat 1876: CCXXVII). Puerto Rico.
- E. scalaris* (Boheman 1840: 349) [= *E. hieroglyphicus* (Sturm 1826: 115), *nom. nud.*]. Costa
Rica and Cuba.
- E. scalptus* Champion 1911: 259. Costa Rica.
- E. similis* (Drury 1773: pl. XXXIII) [= *E. rufescens* (Drury 1773: pl. XXXIII); *E. cameleon*
(Olivier 1790: 543); *E. amabilis* (Waterhouse 1879: 309); *E. bicolor* (Cockerell 1893: 375);
E. concolor (Cockerell 1893: 375)]. Jamaica.
- E. sphacelatus* (Olivier 1807: 319). Cuba (?), Dominican Republic, Haiti, Jamaica, and Puerto
Rico.
- E. squamipennis* Gemminger 1871: 119 [= *E. lepidopterus* (Gyllenhal 1840: 353), not Gyllenhal
1834]. Hispaniola.
- E. sulcicrus* Champion 1911: 268. Colombia, Costa Rica, Guatemala, Nicaragua, and Panama.
- E. sulcipennis* Champion 1911: 265. Guatemala.
- E. sulphuratus* Chevrolat 1879: XC VII. Cuba.
- E. tessellatus* Champion 1911: 262. Costa Rica.
- E. triangulifer* Champion 1911: 261. Costa Rica and Panama.
- E. verecundus* (Chevrolat 1833: 16) [= *E. modestus* (Gyllenhal 1840: 345)]. El Salvador and
Mexico.
- E. vermiculatus* Champion 1911: 264. Mexico.
- E. viridans* (Boheman 1840: 379). St. Thomas.
- E. viridilineatus* Champion 1911: 266. Panama.
- E. viridipupillatus* (Cockerell 1893: 374). Jamaica.
- E. vitraci* (Fleutiaux & Sallé 1889: 380). Guadeloupe.
- E. vittatus* (Linnaeus 1758: 380) [= *E. pulcher* (Brown 1776: 126); *E. inornatus* (Cockerell
1893: 375); *E. musae* (Cockerell 1893: 374); *E. rubrovittatus* (Cockerell 1893: 375); *E.
virescens* (Cockerell 1893: 375); *E. jamaicensis* Bovie 1907: 69]. Jamaica.

E. vitticollis Champion 1911: 256 [= *E. caerulescens* Champion 1911: 256, **syn. nov.**; *E. mutatus* Champion 1911: 256, **syn. nov.**]. Belize and Guatemala.

- Exorides* Pascoe 1881: 43
E. abruptecostatus Marshall 1922: 210. Ecuador.
E. bifurcatus Marshall 1926: 537. Colombia.
E. caudatus Marshall 1922: 217 [= *E. rufelii* Voss 1953: 59]. Colombia.
E. corrugatus Marshall 1922: 218. Venezuela.
E. cylindricus Marshall 1922: 213. Colombia.
E. equatorius Marshall 1922: 219. Ecuador and Peru.
E. equicaudatus Marshall 1922: 207. Ecuador.
E. espeletiae (Kirsch 1889: 23) [= *E. espeletiae* Lona 1938: 525, lapsus]. Colombia.
E. festivus Marshall 1952: 325. Venezuela.
E. immadidus Kuschel 1950: 71. Peru.
E. labyrinthicus (Kirsch 1889: 22). Colombia.
E. lajoyei (Bovie 1907: 68). Venezuela.
E. lindingi (Kirsch 1889: 24). Colombia.
E. marshalli (Bovie 1908: 44). Colombia.
E. masoni Marshall 1922: 215. Colombia.
E. mucronatus (Faust 1892: 18) [= *E. inflatus* Marshall 1922: 214]. Colombia and Venezuela.
E. obesus Marshall 1922: 209. Colombia.
E. octocostatus Hustache 1938: 117. Peru.
E. otiosus Hustache 1938: 116. Ecuador.
E. pedunculatus Marshall 1922: 216. Ecuador.
E. praeteritus Marshall 1922: 206 [= *E. carinatus* Olliff 1891: 65, misident., not Pascoe, 1881].
 Ecuador.
E. pyriformis Marshall 1922: 211. Colombia.
E. quadriovittatus (Kirsch 1889: 23). Colombia.
E. rugosus (Taschenberg 1870: 188). Colombia.
E. septemcostatus Marshall 1922: 212. Colombia and Ecuador.
E. sulcicollis (Kirsch 1889: 24). Ecuador.
E. wagneri (Harold 1863: 176) [= *E. carinatus* Pascoe 1881: 43]. Ecuador.
E. whymperi (Olliff 1891: 63). Ecuador.
E. willei Kuschel 1950: 72. Peru.

Hormotrophus Schoenherr 1843: 43

H. aureomixtus Boheman 1843: 44. Hispaniola.

Oxyderces Schoenherr 1823: col. 1140 [= *Placocompsus* Marshall 1922: 201]

- O. argentinicus* (Heller 1921: 28). Argentina.
O. bimaculatus (Hustache 1923: 290). Argentina and Bolivia.
O. cinereus (Hustache 1938: 110). Ecuador.
O. cretaceus (Fabricius 1792: 452) [= *O. scutellaris* (Chevrolat 1880: 197)]. Guadeloupe and Martinique.
O. croesus (Faust 1892: 11). Venezuela.
O. dubius (Voss 1932: 40). Paraguay.
O. euchloris (Pascoe 1880: 424). Peru.
O. exaratus (Hustache 1938: 111). Colombia.

- O. mansuetus* (Hustache 1938: 109). Colombia and Ecuador.
- O. mirandus* (Pascoe 1880: 423) [= *E. cicatricosus* (Hustache 1938: 105); *E. nigroundulatus* (Hustache 1938: 104); *E. texatus* (Hustache 1938: 114)]. Colombia and Ecuador.
- O. sinuatocostatus* (Hustache 1938: 112). Brazil.
- O. tarapotae* (Hustache 1938: 113). Peru.
- O. viridiaceris* (Hustache 1938: 115). Colombia.
- O. viridipes* (Bohemian 1840: 179). Colombia.
- Parthenides* Kuschel 1986: 79 [= *Parthenius* Kuschel 1955: 306, not Gistel 1857]
- P. virginalis* (Olivier 1807: 341). Brazil.
- Phaops* Schoenherr 1823: col. 1140 [= *Phaops* Sahlberg 1823: 25, not available, type species not fixed; *Eustales* Germar 1824: 443; *Eustalis* Germar 1824: 443, incorrect original spelling, emended Schoenherr 1826; *Emmeria* Pascoe 1881: 42]
- P. adamantinus* (Germar 1824: 444) [= *P. scintillans* (Rosenschoeld 1840: 363)]. Brazil.
- P. albomarginatus* (Rosenschoeld 1840: 368). Brazil.
- P. albugo* Sahlberg 1823: 27 [= *P. leucogaeus* (Germar 1824: 445)]. Brazil and Hispaniola.
- P. ambitiosus* (Bohemian 1840: 373) [= *P. aegrotus* (Bohemian 1840: 371)]. Argentina, Brazil, and Paraguay.
- P. angustifrons* (Lucas 1857: 159). Peru.
- P. cerussatus* (Bohemian 1833: 651) [= *P. micans* (Kirsch 1874: 389)]. Bolivia, Brazil, and Peru.
- P. curvimanus* (Champion 1911: 272). Guatemala.
- P. harmonicus* (Germar 1824: 445). Brazil.
- P. marginatus* (Pascoe 1881: 43). Brazil.
- P. opalinus* (Bohemian 1833: 652). Brazil and French Guiana.
- P. peruvianus* (Hustache 1938: 278). Bolivia and Peru.
- P. thunbergii* (Dalman 1823: 84) [= *P. pentachordius* (Germar 1824: 446)]. Brazil.
- P. vestalis* (Pascoe 1880: 423) [= *P. puncticollis* (Hustache 1940: 273); *P. versicolor* Kuschel 1955: 308, nom. nud.]. Argentina, Bolivia, Brazil, Ecuador, French Guiana, and Peru.
- Phaopsis* Kuschel 1955: 306
- P. laeta* (Bohemian 1840: 374) [= *P. virgata* (Bohemian 1840: 375)]. Argentina, Brazil, and Paraguay.
- P. thalassina* (Bohemian 1840: 378). Brazil.
- Platyomus* Sahlberg 1823: 29 [= *Platyomus* Schoenherr 1823: col. 1140, not Sahlberg 1823]
- P. auromaculatus* (Voss 1932: 37). Costa Rica.
- P. chrysopus* (Champion 1911: 286). Belize, Guatemala, and Mexico.
- P. geminus* (Champion 1911: 284). Mexico.
- P. macroscapus* (Champion 1911: 285). El Salvador, Guatemala, and Honduras.
- P. nigroguttatus* (Champion 1911: 284). Mexico.
- P. squaminatus* (Champion 1925: 429). Costa Rica.
- P. zebra* (Champion 1911: 286). Mexico.
- Pseudeustylus* Champion 1911: 296
- P. cupreotinctus* Champion 1911: 297. Guatemala.
- P. cupreoviridis* Champion 1911: 297. Costa Rica and Panama.

P. planicollis Marshall 1934: 626. Cuba.

Pseudocyphus Schaeffer 1905: 179

P. eximus Voss 1932: 36. Costa Rica.

Rhinospathe Chevrolat 1878: XIX

R. albomarginata Chevrolat 1878: XX. Costa Rica and Panama.

R. v-album Campion 1911: 249. Guatemala.

Simophorus Faust 1892: 19

S. pustulosus Faust 1892: 19. Venezuela.

Synthlibonotus Schoenherr 1847: 41

S. rufipes Schoenherr 1847: 41 [= *S. rufipes* Faust 1892: 17, not Schoenherr 1847]. Colombia and Venezuela.

S. scapha Faust 1892: 18. Venezuela.

S. viator Chevrolat 1880: XLII. Guatemala and Mexico.

Xestogaster Marshall 1922: 22

X. mucorea (Kirsch 1889: 17). Peru.

X. porosa Marshall 1922: 221. Colombia.

X. squalida Marshall 1922: 222. Colombia.

X. viridilimbata (Bovie 1907: 67). Brazil.

Subtribe OTIORHYNCHINA Schoenherr 1823

Oosomina Schoenherr 1823.

Peritelina Lacordaire 1863.

Trachyphloeina Lacordaire 1863.

Brachyrhinina Bedel 1885.

Epitosus Sharp 1891: 150 (**new placement**, R. Anderson, personal comm.)

E. boops Sharp 1891: 150. Guatemala.

Eucyllus Horn, 1876: 74

E. echinus Van Dyke 1936: 31. *Mexico.

Geoderodes Casey 1888: 265

G. hispidus Horn 1894: 442. Mexico.

Orthoptochus Casey 1888: 261

O. squamiger Casey 1888: 261. *Mexico.

Otiorhynchus Germar 1824: 343 [= *Brachyrhinus* Latreille 1802: 200, rejected genus-group name; *Panaphilis* Dejean 1821: 92, rejected genus-group name; *Loborrhynchus* Schoenherr 1823: col. 1145, rejected genus-group name; *Otiorrhynchus* Gemminger & Harold 1871: 2250, lapsus]

O. cribricollis Gyllenhal 1834: 582. Europe. Introduced into Mexico and the United States.

- O. juvencus* Gyllenhal 1834: 609 [= *O. deustus* Blanchard 1851: 381]. Chile (introduced from Europe).
O. rugosostriatus (Goeze 1777: 395 [= *O. scabrosus* (Marsham 1802: 298)]. Europe. Introduced into Canada, Chile, and the United States.
O. sulcatus (Fabricius 1775: 155). Europe. Introduced into Canada, Chile, and the United States.

Peribrotus Gerstaecker 1871: 72 [= *Euperitelus* Champion 1911: 341]
P. albovarius (Champion 1911: 341). Mexico.

Sciorhinus Sharp 1891: 142 (**new placement**, R. Anderson, personal comm.)
S. pictus Sharp 1891: 142. Mexico.

Thricolepis Horn 1876: 68
T. inornata Horn 1876: 69. *Mexico.
T. seminuda Horn 1894: 443. Mexico.

Trachyphloeomimus Champion 1911: 342 [= *Thamiras* Champion 1911: 343]
T. alternatus O'Brien 1972: 177. Mexico.
T. championi O'Brien 1972: 170. Mexico.
T. mexicanus O'Brien 1972: 175. Mexico.
T. sharpi O'Brien 1972: 174. Mexico.
T. solitarius (Sharp 1911: 177). Mexico.
T. spurcus Champion 1911: 342. Mexico.
T. undulatoides O'Brien 1972: 176. Mexico.
T. undulatus (Champion 1911: 343). Guatemala.

Subtribe POLYDROSINA Schoenherr 1823

Brachyderina Schoenherr 1826.
Cneorhinina Lacordaire 1863.
Scythropina Lacordaire 1863.

Ameladus Sharp 1911: 171
A. inornatus Sharp 1911: 171. Mexico.

Apodrusus Marshall 1922: 59 [= *Apodrusus* Wolcott 1924: 130, lapsus]
A. argentatus Wolcott 1924: 130. Mona and Puerto Rico.
A. wolcotti Marshall 1922: 59. Puerto Rico.

Cautoderus Sharp 1911: 169
C. mexicanus Sharp 1911: 170. Mexico.
C. nigrocinctus Champion 1911: 339. Mexico and Nicaragua.

Chaetopantus Sharp 1911: 170
C. illustris Sharp 1911: 170. Mexico.

Mestorus Schoenherr 1840: 910
M. adumbratus Fahraeus 1840: 911. Mexico.
M. crinitus Sharp 1891: 148. Mexico.
M. gracilis Fahraeus 1840: 911. Mexico.

Mitostylus Horn 1876: 107 [= *Derosomus* Sharp 1891: 168]
M. fragilis (Sharp 1891: 168). Mexico.
M. glaucus (Champion 1911: 339). Mexico.
M. gracilis Horn 1894: 444. Mexico.
M. scutellaris (Sharp 1911: 169). Guatemala, Honduras, and Mexico.
M. setosus (Sharp 1891: 168). *Mexico.
M. tenuis Horn 1876: 107. *Mexico.

Polydrosodes Champion 1911: 214
P conicus Champion 1911: 215. Mexico.

Polydrusus Germar 1817: 341 [= *Phyllerastes* Schoenherr 1823: col. 1141; *Polydrosus* Schoenherr 1823: col. 1144; *Polydrosus* Schoenherr 1826: 138; *Eustolus* Thomson 1859: 131; *Cyphomimus* Horn 1876: 105; *Thomsoneonymus* Desbrochers 1902: 113]
P. acuminatus (Champion 1911: 213). Mexico.
P. amplipennis (Champion 1911: 213). Mexico.
P. chilensis Kuschel 1950: 182. Chile.
P. chlorogaster (Champion 1911: 213). Guatemala.
P. delicatulus (Horn 1894: 444). Mexico.
P. flavonotatus (Champion 1911: 210). Guatemala.
P. fuscofasciatus (Champion 1911: 210). Guatemala, Honduras, Mexico, and Nicaragua.
P. guadelupensis (Hustache 1929: 186). Guadeloupe.
P. immaculatus (Champion 1911: 212). Guatemala.
P. latitarsis (Hustache 1929: 185). Guadeloupe.
P. longicornis (Champion 1911: 210). Guatemala and Honduras.
P. macrocephalus (Champion 1911: 212). Mexico.
P. mutabilis (Champion 1911: 211). Costa Rica and Panama.
P. nothofagi Kuschel 1950: 183. Argentina and Chile.
P. pallidisetis (Champion 1911: 214). Mexico.
P. partitus (Champion 1911: 211). Mexico.
P. peninsularis (Horn 1894: 445). Mexico.
P. roseus (Blanchard 1851: 374) [= *P. robigo* Kuschel 1950: 184]. Argentina and Chile.

Pseudopantomorus Champion 1911: 338
P. latifrons Champion 1911: 338. Panama.

Pythis Champion 1911: 340
P. amplicollis Champion 1911: 340. Mexico.

Sciadrusus Sharp 1911: 171
S. propheticus Sharp 1911: 172. Mexico.

Sciomias Sharp 1911: 172
S. elegans Sharp 1911: 172. Mexico.
S. latipennis Sharp 1911: 174. Guatemala.
S. rugicollis Sharp 1911: 173. Guatemala.
S. subtilis Sharp 1911: 173. Mexico.

Polydrosina incertae sedis
Curculio chloroticus Olivier 1807: 348. Antilles.
Curculio fulgens Olivier 1807: 329. Puerto Rico.
Curculio religiosus Olivier 1807: 302. Guadeloupe.

Subtribe ANYPOTACTINA Champion 1911

Anypotactus Schoenherr 1840: 299 [= *Anisactus* Kuschel 1955:275; *Parasomus* Sharp 1911: 174]
A. bicaudatus Champion 1911: 215. Costa Rica and Panama.
A. exilis Boheman 1840: 300. Colombia, Costa Rica, Guatemala, Honduras, Nicaragua, Panama, and Venezuela.
A. gracilis Voss 1932: 35 [= *A. curvipes* Hustache 1940: 272; *A. peruvianus* Hustache 1938: 265]. Bolivia and Peru.
A. jansoni (Sharp 1911: 174). Costa Rica, El Salvador, Honduras, and Nicaragua.
A. morosus (Boheman 1840: 449). Colombia.
A. strangulatus Hustache 1938: 266. Bolivia and Peru.
A. sulcicollis Faust 1892: 21. Venezuela.
Bothinodontes Kirsch 1868: 241 [= *Bothynodontes* Gemminger & Harold 1871: 2242, unjust. emend.]
B. gibbipennis (Champion 1911: 332). Colombia and Mexico.
B. squalidus Kirsch 1868: 243. Colombia.

Cylloproctus Faust 1892: 22
C. dissimilis Voss 1936: 105. Brazil.
C. modestus Faust 1892: 23. Colombia.
C. murinus Faust 1892: 23. Venezuela.
C. nigropictus Voss 1936: 108. Brazil.
C. pantomoroides Voss 1936: 109. Colombia.
C. pustulatus Voss 1936: 107. Brazil.
C. pyriformis Faust 1892: 24. Venezuela.
C. simoni Faust 1892: 25. Venezuela.
C. vehemens Faust 1892: 24. Venezuela.

Helicorrhynchus Olliff 1891: 61 [= *Helicorrhynchus* Lona 1937: 240, lapsus]
H. vulsus Olliff 1891: 62. Ecuador.

Hyphantus Germar 1824: 334
H. angulatus Vaurie 1963: 282. Brazil.
H. argentinensis Hustache 1926: 156. Argentina, Brazil, Paraguay, and Uruguay.

- H. baccifer* Germar 1824: 335. Brazil.
H. bondari Vaurie 1963: 290. Brazil.
H. bracteatus Vaurie 1963: 272. Brazil.
H. brevicauda Vaurie 1963: 266. Brazil.
H. carinatus Vaurie 1963: 263. Brazil.
H. chryseus Vaurie 1963: 278. Brazil.
H. conicus Vaurie 1963: 277. Brazil.
H. dehiscens Desbrochers 1892: 18. Brazil.
H. dentipes Soares & Scivittaro 1977: 35. Brazil.
H. desbrochersi Soares & Scivittaro 1977: 36. Brazil.
H. diringshofeni Soares & Scivittaro 1977: 36. Brazil.
H. distinguendus Desbrochers 1892: 17. Brazil.
H. fausti Vaurie 1964: 37. Brazil.
H. gibbosus Vaurie 1963: 280. Brazil.
H. gnomus Vaurie 1963: 274. Brazil.
H. hustachei Vaurie 1963: 284 [= *H. granulicollis* Jacob 1937: 156, nom. nud.]. Argentina,
Brazil, and Paraguay.
H. hypercalus Vaurie 1963: 271. Brazil.
H. ignotus Soares & Scivittaro 1977: 39. Brazil.
H. incongruus Vaurie 1963: 267. Brazil.
H. inoratus Soares & Scivittaro 1977: 41. Brazil.
H. longicauda Vaurie 1963: 295 [= *H. lanceolatus* Vaurie 1963: 297]. Brazil.
H. maculifer Desbrochers 1892: 16. Brazil.
H. matronalis Vaurie 1963: 280. Brazil.
H. minutus Vaurie 1963: 288. Argentina and Brazil.
H. montanus Vaurie 1963: 260. Brazil.
H. olivae Vaurie 1963: 298. Brazil.
H. parvulus Vaurie 1963: 287. Brazil.
H. pyramis Vaurie 1963: 275. Brazil.
H. quadridens Vaurie 1971: 123. Brazil.
H. serpenis Vaurie 1963: 262. Brazil.
H. setulosus (Bohemian 1859: 127). Argentina and Uruguay.
H. simulans Vaurie 1963: 294 [= *H. sulcifrons* Hustache 1926: 156, misident., not Boheman
1843]. Argentina, Brazil, and Uruguay.
H. squamosus Vaurie 1963: 291. Brazil.
H. subminutus Vaurie 1963: 286. Argentina and Brazil.
H. sulcatus Soares & Scivittaro 1977: 44. Brazil.
H. sulcifrons Boheman 1843: 410. Argentina, Brazil, and Uruguay.
H. teretirostris Desbrochers 1892: 19. Brazil.
H. tibialis Vaurie 1963: 281. Brazil.
H. titan Vaurie 1963: 270. Brazil.
H. uncinatus Desbrochers 1892: 18. Brazil.
H. vauriaeae Soares & Scivittaro 1977: 45. Brazil.
H. verrucifer Boheman 1843: 409. Brazil.
H. zikani Soares & Scivittaro 1977: 46. Brazil.

Hypsometopus Kirsch 1868: 222

H. inquinatus Kirsch 1868: 223. Colombia.

Neoanyptactus Hustache 1938: 266

N. brevirostris (Hustache 1938: 266). Bolivia.

N. marshalli Kuschel 1950: 69. Peru.

Nototactus Kuschel 1952: 231

N. angustirostris Kuschel 1952: 233. Chile.

N. latirostris Kuschel 1952: 232 [= *N. wittmerianus* Kuschel 1952: 233]. Argentina and Chile.

Phanasora Pascoe 1881: 38 [= *Baccharisphila* Hustache 1939: 66]

P. benoisti (Hustache 1939: 67). Ecuador.

P. plumbea Pascoe 1881: 39. Colombia.

Polydacrys Schoenherr 1834: 130 [= *Polydacris* Lacordaire 1863: 66, lapsus]

P. brevicollis Champion 1911: 220. Guatemala.

P. depressifrons Boheman 1840: 298 [= *P. nubilosus* (Bohemian 1840: 296); *P. cavirostris* (Schaeffer 1908: 214)]. *Belize, Costa Rica, El Salvador, Grenada, Guatemala, Honduras, Mexico, Nicaragua, Panama, Puerto Rico, and St. Vincent.

P. moestus Chevrolat 1880: 190 [= *P. nigrosparsus* Chevrolat 1880: 190]. Guadeloupe.

P. mucronatus Champion 1911: 219. Costa Rica and Panama.

P. nigrofasciatus Champion 1911: 219. El Salvador and Mexico.

P. scansorius (Klug 1829: 13) [= *P. modestus* Gyllenhal 1834: 131; *P. luctuosus* Dejean 1837: 278, nom. nud.]. Cuba, Dominican Republic, and Guadeloupe.

P. seriegranosus Champion 1911: 218. Bolivia, Colombia, and Panama.

Prepodellus Kirsch 1868: 239

P. nigriclavis Kirsch 1868: 240. Colombia.

P. ruficornis Kirsch 1868: 240. Colombia.

P. viridisquamis Champion 1911: 220. Costa Rica, Mexico, Nicaragua, and Panama.

Subtribe BARYNOTINA Schoenherr 1826

Anomadus Horn 1876: 21

A. obliquus Horn 1876: 21. Mexico.

Apotomoderes Dejean 1834: 253 [= *Apotomus* Schoenherr 1834: 44; *Eurilia* Laporte 1840: 308]

A. lateralis (Gyllenhal 1834: 45). Hispaniola.

Bradyrhynchoides Pierce 1913: 412

B. rugicollis (Sharp 1891: 142). Mexico.

Buromicrus Sharp 1891: 145

B. cristatus Sharp 1891: 146. Guatemala and Nicaragua.

B. globipennis Sharp 1891: 146. Guatemala and Honduras.

B. marshalli Sleeper 1959: 108. Guatemala.

B. squamosus Sharp 1891: 145. Belize, Guatemala, and Honduras.

Chauliopleurus Champion 1911: 270

C. adipatus Champion 1911: 270. Costa Rica.

C. quadrifoveatus Champion 1911: 271. Costa Rica.

C. rufovittatus Champion 1911: 271. Costa Rica.

C. viridisquamans Champion 1925: 428. Costa Rica.

Claeoteges Pascoe 1880: 427 [= *Cloeoteges* Blackwelder 1947: 797, lapsus]

C. granulosa Champion 1911: 276. Belize, Guatemala, Honduras, and Mexico.

C. oblitterata Champion 1911: 277. Guatemala, Honduras, and Mexico.

C. tuberculosa Champion 1911: 277. Guatemala.

C. virosa Pascoe 1880: 428. Costa Rica, Guatemala, Nicaragua, and Panama.

Cleistolophus Sharp 1891: 143

C. instabilis Sharp 1891: 144. Guatemala.

C. similis (Chevrolat 1880: LXI). Belize, Costa Rica, Guatemala, Honduras, Mexico, and Nicaragua.

C. subfasciatus Sharp 1891: 143. Guatemala, Honduras, Mexico, and Nicaragua.

C. viridimargo Champion 1911: 331. Belize and Honduras.

Cryptolepidus Van Dyke 1936: 191 [= *Lepidopus* Van Dyke 1936: 76, not Gouan 1770;

Pseudoeucylus Tanner 1950: 71]

C. cazieri (Van Dyke 1936: 73). *Mexico.

Decasticha Champion 1911: 273

D. decemguttata Champion 1911: 275. Costa Rica.

D. quadriplagiata Champion 1911: 274. Panama.

D. subocellata Champion 1911: 273. Costa Rica.

D. tetrastigma Champion 1911: 274. Costa Rica.

D. tricostata Champion 1925: 429. Panama.

Doleropus Buchanan 1947: 46

D. apterus Buchanan 1947: 47. Cuba.

Epicaerus Schoenherr 1834: 323 [= *Epagrius* Schoenherr 1840: 119; *Bradyrhynchus* Sharp

1891: 140; *Cacochromus* Sharp 1891: 140; *Diorynotus* Sharp 1891: 105; *Epagrius* Sharp

1891: 128, misident., not Schoenherr 1840; *Epiacaerus* Horn 1894: 357, lapsus; *Melbonus*

Casey 1895: 820; *Epagriopsis* Champion 1911: 328; *Epicoerus* Champion 1912: 182, lapsus]

E. aequalis Sharp 1891: 107. Mexico.

E. albosquamatus (Chevrolat 1880: XLIII). Costa Rica and Mexico.

E. amulae Sharp 1891: 120. Mexico.

E. aurifer Boheman 1842: 278 [= *E. ravidus* Boheman 1842: 277]. Mexico.

E. bicolor Sharp 1891: 116. Guatemala and Mexico.

E. biformis Sharp 1891: 127. Guatemala.

E. brevirostris (Sharp 1891: 141). Mexico.

E. calvus Sharp 1891: 106. Mexico.

- E. capetillensis* Sharp 1891: 115. Belize and Guatemala.
E. carinatus Boheman 1842: 285. Guatemala and Mexico.
E. carteri Chevrolat 1880: LXII. Costa Rica and Guatemala.
E. centralis Sharp 1891: 107. Mexico.
E. championi Sharp 1891: 114. Guatemala and Mexico.
E. cognatus Sharp 1891: 105. Mexico.
E. concolor Sharp 1891: 126. Mexico.
E. constans (Sharp 1891: 134). Guatemala.
E. convexus Boheman 1842: 281. El Salvador, Guatemala, and Mexico.
E. costatus Sharp 1891: 109. Mexico.
E. costicollis Sharp 1891: 109. Mexico.
E. coxalis Sharp 1891: 108. Mexico.
E. cultripennis Boheman 1842: 273. Mexico.
E. curvipes (Sharp 1891: 131). Mexico.
E. decoratus Sharp 1891: 121. Mexico.
E. depilis Boheman 1842: 279. Mexico.
E. durangoensis Sharp 1891: 119. Mexico.
E. elegantulus Champion 1911: 326. Mexico.
E. fallax Boheman 1842: 274. Mexico.
E. foveicollis (Sharp 1891: 135). Mexico.
E. foveifrons (Champion 1911: 327). Mexico.
E. fronterae Sharp 1891: 112. Mexico.
E. godmani Sharp 1891: 124. Mexico.
E. grandis (Sharp 1891: 140). Mexico.
E. griseus Boheman 1842: 282 [= *E. sphaeroides* Boheman 1842: 283]. Mexico.
E. hispidus (Sharp 1891: 133). Mexico.
E. hoegei Sharp 1891: 125. Mexico.
E. hystriculus (Sharp 1891: 133). Mexico.
E. imbricatus (Say 1824: 317). *Mexico.
E. impar Sharp 1891: 128. Guatemala.
E. inaequalis (Sharp 1891: 139). Costa Rica, Guatemala, Honduras, and Panama.
E. inflatus Sharp 1891: 112. Mexico.
E. jugicola (Sharp 1891: 135). Guatemala.
E. laevinasus (Sharp 1891: 138). Guatemala.
E. lateralis Sharp 1891: 120. Mexico.
E. lucanus Horn 1894: 441. Mexico.
E. macropterus Sharp 1911: 324. Mexico.
E. macropus Champion 1911: 324. Mexico.
E. marginatus Sharp 1891: 122 [= *E. insolitus* Sharp 1891: 122]. Mexico.
E. mexicanus Boheman 1834: 324 [= *E. aeruginosus* Boheman 1842: 276; *E. cyphus* Boheman 1842: 272; *E. transversepunctatus* Boheman 1842: 271]. *Mexico.
E. minor Sharp 1891: 118. Guatemala.
E. monclovae Sharp 1891: 118. Mexico.
E. nebulosus Boheman 1834: 326 [= *E. neglectus* Boheman 1834: 325]. Mexico.
E. niger Sharp 1891: 106. Mexico.
E. nubilosus (Bohemian 1840: 121) [= *E. metallescens* (Bohemian 1840: 122); *E. tolucae* (Sharp 1891: 141)]. Mexico.

- E. oculatus* Sharp 1891: 121. Mexico.
E. opacus (Sharp 1891: 135). Guatemala and Mexico.
E. opercularis (Say 1831: 9) [= *E. gravidus* (Sharp 1891: 130); *E. morosus* (Sharp 1891: 131)].
 Mexico.
E. oscillator Sharp 1891: 117. Mexico.
E. paradae (Sharp 1891: 132). Mexico.
E. pavidus Sharp 1891: 127. Guatemala.
E. pedestris Sharp 1891: 115. Guatemala.
E. planirostris Sharp 1891: 119. Mexico.
E. praeteritus (Sharp 1891: 137). Mexico.
E. pumilus (Sharp 1891: 138). Guatemala.
E. pyriformis Sharp 1891: 124. Mexico.
E. retrorsus (Champion 1911: 330). Mexico.
E. reversus Sharp 1891: 113. Mexico.
E. salvadorensis Kuschel 1956: 333. El Salvador.
E. samson (Sharp 1891: 139). Belize, Costa Rica, Guatemala, and Mexico.
E. scutellaris Sharp 1891: 123. Mexico.
E. sexcostatus Sharp 1891: 125. Mexico.
E. simplex (Sharp 1891: 134). Guatemala and Mexico.
E. smithi (Sharp 1891: 136). Mexico.
E. squalidus Sharp 1891: 116. Mexico.
E. squamosus Sharp 1891: 122. Mexico.
E. sturmi Sharp 1891: 114. Mexico.
E. sulcirostris Sharp 1891: 106. Mexico.
E. tenuis Sharp 1891: 115. Mexico.
E. tristis (Chevrolat 1880: XLII). Mexico.
E. truquianus Champion 1911: 327. Mexico.
E. variolosus (Sharp 1891: 132). Mexico.

Eugeonemus Buchanan 1947: 44 [= *Geonemus* of authors, not Schoenherr 1834]

- E. alternans* (Klug 1829: 13) [= *E. alternans* (Boheman 1834: 293)]. Cuba.
E. aureosquamatus (Jacquelin du Val 1857: 195). Cuba.
E. irregularis (Perroud 1853: 518). Cuba.
E. robustus (Perroud 1853: 515). Cuba.

Eumestorus Sharp 1891: 148

- E. luctuosus* (Chevrolat 1880: LXII). Belize and Mexico.
E. proximus Sharp 1891: 147. Mexico.

Ischnioplus Chevrolat 1878: IX

- I. niveoguttatus* Chevrolat 1878: X. Cuba.
I. viridiguttatus Chevrolat 1878: X. Dominican Republic.

Lachnopus Schoenherr 1840: 380 [= *Menoetius* Dejean 1821: 94, rejected genus-group name;
Ptilopus Schoenherr 1823: col. 1140, rejected genus-group name]

- L. acuticollis* (Gyllenhal 1834: 37). Cuba.
L. aereus (Gyllenhal 1834: 40) [= *L. blandus* (Gyllenhal 1834: 26)]. Hispaniola.
L. alboguttatus Marshall 1934: 622. Cuba.

- L. albomaculatus* (Gyllenhal 1834: 37). Hispaniola.
- L. argus* (Reiche 1840: 275). *Cuba.
- L. atramentarius* (Gyllenhal 1834: 33). Dominican Republic and Haiti.
- L. aulicus* (Gyllenhal 1834: 35). Hispaniola.
- L. aurifer* (Drury 1773: 68). Cuba and Jamaica.
- L. bellus* Marshall 1926: 54. Haiti.
- L. bivirgatus* Marshall 1934: 621. Cuba.
- L. bruneri* Marshall 1933: 59. Cuba.
- L. buchanani* Marshall 1933: 59. Cuba.
- L. campechianus* Gyllenhal 1840: 388 [= *L. adspersus* Chevrolat 1880: 191]. Guadeloupe.
- L. canescens* Gyllenhal 1840: 388. Hispaniola.
- L. chirographus* (Olivier 1807: 334). Hispaniola.
- L. chlorophanus* (Gyllenhal 1834: 39). Haiti.
- L. coffeeae* Marshall 1922: 60 [= *L. montanus* Marshall 1922: 61, **syn. nov.**]. Puerto Rico.
- L. consentaneus* Perroud 1853: 487. Hispaniola.
- L. curvipes* (Fabricius 1787: 113) [= *L. calcaratus* (Olivier 1807: 350), **syn. nov.**]. Dominican Republic, Guadeloupe, Jamaica, Nevis, Puerto Rico, St. Barthélemy, St. Croix, St. Kitts, St. Thomas, St. Vincent, and Tortola.
- L. dentipes* Perroud 1853: 489. Hispaniola.
- L. distortus* Gyllenhal 1840: 393 [= *L. valgus* (Herbst 1795: 129), misident., not Fabricius 1775]. West Indies.
- L. gowdeyi* Marshall 1926: 531. Jamaica.
- L. granicollis* Gyllenhal 1840: 390. Hispaniola.
- L. guerinii* Jacquelin du Val 1857: 185. Cuba.
- L. hirtus* Perroud 1853: 484. Hispaniola.
- L. hispidus* (Gyllenhal 1834: 34). *Cuba.
- L. histrio* Marshall 1926: 534 [= *L. erectosetosus* Marshall 1926: 536, **syn. nov.**]. Haiti.
- L. inconditus* Rosenschöeld 1840: 383. Dominican Republic and Haiti.
- L. interruptus* Perroud 1853: 475. Hispaniola.
- L. kofyesi* Wolcott 1941: 104. Mona.
- L. lineatoguttatus* Perroud 1853: 468 [= *L. seriepunctatus* Jacquelin du Val 1857: 187]. Cuba.
- L. lineicollis* (Chevrolat 1880: 175) [= *L. foiveicollis* (Chevrolat 1880: 175); *L. quadrataenia* (Chevrolat 1880: 190)]. Guadeloupe.
- L. luctuosus* (Klug 1829: 13) [= *L. luctuosus* (Gyllenhal 1834: 26), not Klug 1829]. Cuba.
- L. luxurians* (Olivier 1807: 334). Hispaniola.
- L. memnonius* (Gyllenhal 1834: 42). St. Barthélemy.
- L. mercator* (Olivier 1807: 335). Hispaniola.
- L. multipunctatus* Jacquelin du Val 1857: 190. Cuba.
- L. mundus* Gyllenhal 1834: 31. Dominican Republic.
- L. niveoirroratus* Jacquelin du Val 1857: 189. Cuba.
- L. oteroi* Marshall 1933: 60. Cuba.
- L. planifrons* Gyllenhal 1840: 385. Dominican Republic.
- L. plebejus* (Gyllenhal 1834: 23). Hispaniola.
- L. plumipes* Perroud 1853: 471 [= *L. wolcotti* Marshall 1926: 533]. Dominican Republic and Haiti.
- L. pollinarius* Gyllenhal 1840: 387. Cuba.
- L. proteus* (Olivier 1807: 336). Haiti.
- L. pruinosus* Gyllenhal 1834: 33. Hispaniola.

L. seini Wolcott 1936: 302. Puerto Rico.
L. sparsimguttatus Perroud 1853: 481. Cuba.
L. splendidus Boheman 1840: 382. Cuba.
L. spretus Gyllenhal 1834: 38. Dominican Republic.
L. sublineatus Perroud 1853: 478 [= *L. sublineatus* Jacquelin du Val 1857: 192, not Perroud 1853]. Cuba.
L. trilineatus Chevrolat 1876: CCXXVIII. Puerto Rico.
L. valgus (Fabricius 1775: 150) [= *L. castilianus* (Herbst 1797: 22)]. Anguilla, Puerto Rico, St. Barthélemy, and St. Croix.
L. villosipes (Boheman 1834: 43). St. Barthélemy and St. Eustatius.
L. vittatus (Klug 1829: 13) [= *L. vittatus* (Gyllenhal 1834: 43)]. Cuba.

Maseorhynchus Sharp 1891: 146
M. hondurensis Sharp 1891: 147. Belize, Guatemala, and Mexico.

Mazenes Champion 1911: 280
M. bifoveatus Champion 1911: 280. Guatemala.
M. geminatus Champion 1911: 281. Guatemala.
M. lineatus Champion 1911: 281. Mexico.

Naupactopsis Champion 1911: 278
N. auropicta Champion 1911: 279. Mexico.
N. delumbis Champion 1911: 278. Mexico.
N. sitonoides Champion 1911: 279. Mexico.

Omileus Horn 1876: 102
O. epicaeroides Horn 1876: 102. *Mexico.

Tetrabothynus Imhoff 1852: no. 68 [= *Tetrabothinus* Lacordaire 1863: 108, lapsus]
T. regalis (Linnaeus 1767: 616). Hispaniola and Peru.
T. spectabilis (Klug 1829: 13) [= *T. spectabilis* (Gyllenhal 1834: 25), not Klug 1829]. Cuba.

Tropirhinus Schoenherr 1823: col. 1140 [= *Callizonus* Schoenherr 1826: 123]
T. elegans (Guérin 1847: 10). Cuba.
T. novemdecimpunctatus (Fabricius 1775: 145) [= *T. olivieri* (Chevrolat 1877: 171)]. French Guiana and Hispaniola.
T. tredecimmaculatus (Guérin 1947: 10). Cuba.

Barynotina incertae sedis
Otiorhynchus cuneiformis Waterhouse 1845: 38. Ecuador (Galapagos Ids.).

Subtribe ANOMOPHTHALMINA Morrone 1998

Anomophthalmus Fairmaire 1884: 498
A. insolitus Fairmaire 1884: 498 [= *A. crawshayi* (Champion 1918: 35)]. Argentina and Chile.
Sysciophthalmus Heller 1906: 5

S. bruchi Heller 1906: 6. Argentina and Chile.

Subtribe TANYMECINA Lacordaire 1863

Piazomiina Reitter 1913.

Pandeliina Pierce 1913.

Airosimus Howden 1966: 174

A. aequatoris Howden 1966: 186. Brazil.

A. amazonensis Howden 1966: 191. Brazil.

A. andersoni Howden 1993: 255. Costa Rica.

A. barbareae Howden 1966: 188. Brazil.

A. boops (Champion 1911: 189). Costa Rica and Panama.

A. callifer (Voss 1934: 69). Costa Rica and Mexico.

A. erubescens (Champion 1911: 188). Costa Rica and Panama.

A. intonsus Howden 1966: 208. Brazil.

A. jacobi (Hustache 1938: 9) [= *A. fasciatus* (Voss 1934: 68); *A. fasciger* (Emden 1939: 257)].
Brazil and Paraguay.

A. kissingeri Howden 1993: 253. Mexico.

A. lucileae Howden 1966: 194. Ecuador.

A. margaritis Howden 1966: 220. Ecuador.

A. merulis Howden 1966: 218. Costa Rica.

A. patienceae Howden 1966: 214. Brazil.

A. pelodus (Warner 1959: 33). Costa Rica.

A. robustus (Faust 1892: 4). Colombia and Venezuela.

A. semirobustus Howden 1966: 199. Colombia.

A. stalagmos Howden 1966: 211. Bolivia and Peru.

A. tuberculifer (Boheman 1840: 295). Brazil.

Aporius Kuschel 1955: 276

A. brevipennis (Chevrolat 1879: CV). French Guiana.

Eurymetopellus Emden 1936: 86 [= *Eurymetopus* Kirsch 1889: 13, misident., not Schoenherr

1840; *Eurymetopinus* Emden 1936: 86, not Winkler 1932]

E. alitcola (Kirsch 1889: 13) [= *E. parvicollis* (Olliff 1891: 67)]. Ecuador.

Galenactus Kuschel 1952: 234

G. litoralis Kuschel 1952: 234. Chile.

Hadromeropsis Pierce 1913: 400 [= *Hadromerus* Schoenherr 1834: 127, not Schoenherr 1823;

Hadromorus Hustache 1926: 157, lapsus; *Hadrorestes* Howden 1982: 85]

H. alacer Howden 1982: 85. Colombia and Ecuador.

H. amoena Howden 1982: 36. Mexico.

H. apicalis Howden 1982: 103. Ecuador.

H. argentinensis (Hustache 1926: 157). Argentina, Brazil, Paraguay, and Uruguay.

H. atomaria (Boheman 1840: 292). Brazil.

H. aurea (Blanchard 1847: 201). Bolivia.

- H. batesi* Howden 1982: 55. Bolivia, Brazil, Costa Rica (?), Panama (?), and Peru.
- H. beverlyae* Howden 1982: 72. Brazil.
- H. bombycina* Howden 1982: 97. Peru.
- H. brachyptera* Howden 1982: 98. Unknown.
- H. brevicoma* Howden 1982: 42. Mexico.
- H. cavifrons* Howden 1982: 122. Peru.
- H. conquisita* Howden 1982: 110. Ecuador.
- H. contracta* Howden 1982: 106. Bolivia.
- H. cretata* (Champion 1911: 184). Costa Rica and Panama.
- H. crinita* Howden 1982: 29. Mexico.
- H. dejeanii* (Bohemian 1840: 293). Mexico.
- H. dialeuca* Howden 1982: 105. Venezuela.
- H. earina* Howden 1982: 124. Ecuador.
- H. excubitor* Howden 1982: 78. Brazil.
- H. exilis* Howden 1982: 92. Bolivia.
- H. fasciata* (Lucas 1857: 156) [= *H. herbacea* (Lucas 1867: 157)]. Brazil.
- H. flagellata* Howden 1982: 30 [= *H. opalina* (Champion 1911: 183, misident., in part)]. Mexico.
- H. fulgens* (Champion 1911: 184). Mexico.
- H. gemmifera* (Bohemian 1845: 418) [= *H. scabricollis* (Faust 1892: 1)]. Colombia, Costa Rica, Guatemala, Panama, and Venezuela.
- H. impressicollis* (Kirsch 1868: 233) [= *H. ruficra* (Kirsch 1868: 232); *H. subaenea* Voss 1953: 61]. Colombia.
- H. inconspecta* Howden 1982: 87. Peru.
- H. institula* Howden 1982: 96. Ecuador.
- H. magica* (Pascoe 1881: 41). Colombia.
- H. mandibularis* Howden 1982: 117. Colombia.
- H. meridiana* Howden 1982: 54. Brazil and Colombia.
- H. micans* (Champion 1911: 183). Guatemala.
- H. nana* Howden 1982: 102. Ecuador.
- H. nebulicola* Howden 1982: 88. Colombia.
- H. nitida* Howden 1982: 115. Ecuador.
- H. nobilitata* (Gyllenhal 1834: 128) [= *H. similis* Hustache 1938: 6]. Argentina and Brazil.
- H. opalina* (Horn 1876: 85) [= *H. viridissima* (Van Dyke 1943: 108)]. *Mexico.
- H. pallida* Howden 1982: 67. Argentina, Brazil, Paraguay, and Uruguay.
- H. pectinata* Howden 1982: 108. Bolivia, Colombia, and Peru.
- H. picchuensis* Howden 1982: 121. Brazil.
- H. plebeia* Howden 1982: 62. Brazil.
- H. pulverulenta* Howden 1982: 65. Brazil.
- H. rufipes* (Champion 1911: 185). Costa Rica.
- H. scamba* Howden 1982: 112. Ecuador.
- H. scintillans* (Champion 1911: 182). Guatemala and Mexico.
- H. silacea* Howden 1982: 90. Colombia.
- H. speculifera* Howden 1982: 69. Brazil.
- H. spiculata* Howden 1982: 119. Bolivia and Peru.
- H. striata* Howden 1982: 126. Colombia.
- H. superba* (Heller 1921: 29). Argentina.

H. togata (Boheman 1840: 290) [= *H. brachispinosa* (Boheman 1840: 291)]. Brazil and Paraguay.

H. transandina Howden 1982: 100. Ecuador.

Isodacrys Sharp 1911: 175

I. apicale Howden 1961: 91. Mexico.

I. brevirostre Howden 1961: 79. Mexico.

I. buchanani Howden 1961: 82. *Mexico.

I. crispum Howden 1961: 84. Mexico.

I. ellipticum Howden 1961: 86. Honduras.

I. geminatum Howden 1961: 80. Mexico.

I. guatemalenum Sharp 1911: 175. Guatemala.

I. mexicanum Sharp 1911: 176. Mexico.

I. minutum Sharp 1911: 176. Mexico.

I. orizabae Sharp 1911: 176. Mexico.

I. ovipennis Schaeffer 1908: 215. *Mexico.

I. schwarzi Champion 1911: 341. Mexico.

Isodrusus Sharp 1911: 207

I. debilis Sharp 1911: 208. *Guatemala and Mexico.

I. guajavus Howden 1970: 50. Jamaica.

I. insulanus Howden 1963: 43. Bahamas.

Macropterus Schoenherr 1840: 419 [= *Megapterum* Gistel 1848: XI]

M. acuminatus Boheman 1840: 421 [= *M. semicostatus* Rosenschoeld 1840: 423]. Brazil.

M. chlorostomus Boheman 1840: 424. Brazil and Colombia.

M. dentipennis Voss 1954: 226. Peru.

M. longipennis Boheman 1840: 420. Brazil.

Minyomerus Horn 1876: 17 [= *Elissa* Casey 1888: 271; *Pseudelissa* Casey 1888: 273]

M. caseyi (Sharp 1891: 151). Mexico.

M. languidus Horn 1876: 18 [= *M. cinereus* (Casey 1888: 274)]. *Mexico.

M. laticeps (Casey 1888: 272). *Mexico.

Pachnaeus Schoenherr 1826: 121 [= *Docorhinus* Schoenherr 1823: col. 1140, *nom. nud.*;

Pachneus Gemminger & Harold 1871: 2224, unjust. emend.]

P. azurescens Gyllenhal 1834: 58 [= *P. griseus* Gyllenhal 1834: 58, **syn. nov.**]. Cuba.

P. citri Marshall 1916: 453. Jamaica.

P. costatus Perroud 1853: 495. Cuba.

P. litus (Germar 1824: 431) [= *P. glaucus* Sturm 1826: 202, *nom. nud.*; *P. opalus* of authors, misident., not Olivier 1807]. *Cuba, Jamaica, and Mexico.

P. marmoratus Marshall 1916: 454. Jamaica.

P. opalus Olivier 1807: 339 [= *P. distans* Horn 1876: 83]. *West Indies.

P. psittacus (Olivier 1807: 339). Cuba and Puerto Rico.

Pandeteinus Champion 1911: 206 [= *Pandeteinus* Guenther & Zumpt 1933: 113, lapsus]

P. elytroplanatus Howden 1959: 378. *Mexico.

P. lucidillus Howden 1959: 378. *Mexico.

- P. magdalenensis* Howden 1959: 375. Mexico.
P. subcancer Howden 1969: 78. Mexico.
P. submetallicus (Schaeffer 1908: 216). *Mexico.
- Pandeleteius* Schoenherr 1834: 129 [= *Pandeletius* Agassiz 1846: 86, unjust. emend.; *Menetypus* Kirsch 1868: 233; *Pandeletius* Gemminger & Harold 1871: 2208, unjust. emend.; *Pandeletejus* Horn 1876: 86, unjust. emend.; *Mentypus* Jacob 1937: 157, lapsus; *Exmenetypus* Voss 1954: 231; *Exmentypes* Warner 1959: 33, lapsus]
- P. admirabilis* Howden 1976: 192. Colombia and Venezuela.
P. albisquamis Champion 1911: 205. Costa Rica, El Salvador, Guatemala, and Mexico.
P. albus (Hustache 1923: 287). Argentina, Bolivia, Brazil, and Paraguay.
P. amulae Champion 1911: 203. Mexico.
P. andeanus Howden 1976: 155. Colombia and Venezuela.
P. angustirostris Faust 1892: 5. Venezuela.
P. antiochensis Howden 1976: 37. Colombia.
P. arcanus Howden 1976: 60. Venezuela.
P. argentatus (Olliff 1891: 62). Ecuador.
P. armatus Champion 1911: 198. Mexico.
P. aspericollis (Hustache 1926: 158). Argentina and Paraguay.
P. assimilis Voss 1934: 69. Brazil.
P. atlas Howden 1976: 113. Venezuela.
P. avilensis Howden 1976: 154. Venezuela.
P. baccatus Howden 1996: 916. Peru.
P. baccharis Kuschel 1949: 16. Chile.
P. bechyneorum Howden 1976: 35. Venezuela.
P. bidentatus Howden 1976: 133. Venezuela.
P. biseriatus (Kirsch 1874: 388) [= *P. rubricatus* (Hustache 1938: 11)]. Argentina, Bolivia, Brazil, Paraguay, and Peru.
P. bordoni Howden 1976: 171. Colombia and Venezuela.
P. brasilianus Voss 1954: 229. Brazil.
P. brevinasus Champion 1911: 200. Guatemala and Mexico.
P. brevipes Champion 1911: 194. Guatemala and Mexico.
P. brevitarsis (Hustache 1938: 10). Argentina.
P. buchanani Howden 1959: 389. *Mexico.
P. campbelli Howden 1976: 104. Colombia.
P. campestris Howden 1976: 73. Colombia and Venezuela.
P. candidus Howden 1976: 187. Venezuela.
P. carinipenis Howden 1976: 123. Colombia.
P. championi Howden 1974: 12. Guatemala and Honduras.
P. chapini Howden 1976: 68. Colombia and Venezuela.
P. ciliatipenis Champion 1911: 204. Mexico.
P. cinereus (Horn 1876: 87). *Mexico.
P. circumcisus Howden 1976: 65. Bolivia.
P. clavisetis Champion 1911: 191. Mexico.
P. clivus Howden 1976: 146. Colombia.
P. colatus Howden 1996: 924. Argentina, Brazil, and Paraguay.
P. commutabilis Howden 1976: 131. Argentina and Bolivia.

- P. confinis* Howden 1996: 885. Bolivia, Brazil, Paraguay, and Peru.
P. confluens Howden 1976: 30. Venezuela.
P. conirostris Howden 1976: 101. Colombia.
P. connatus Howden 1996: 905. Bolivia and Peru.
P. conspersus Champion 1911: 195. Mexico.
P. cornelli Howden 1996: 918. Peru.
P. crispus Champion 1911: 191. Mexico.
P. cucullatus Champion 1911: 192. Mexico.
P. cuneatus Champion 1911: 192. Guatemala, Honduras, and Mexico.
P. cupidus Howden 1976: 174, **stat. nov.** Colombia.
P. defectus Green 1920: 196. *Mexico.
P. dissimilis Voss 1939: 337. Colombia, Costa Rica, Honduras, and Panama.
P. distinctus (Voss 1954: 232). Peru.
P. eberhardi Howden 1976: 135. Colombia.
P. ecuadorensis Howden 1976: 99. Ecuador.
P. ephippiatus Champion 1911: 190. Panama.
P. excisus Howden 1976: 169. Colombia.
P. fasciatus Champion 1911: 201. Guatemala and Mexico.
P. femoralis Champion 1911: 189. Mexico.
P. festivus Howden 1976: 42. Venezuela.
P. flavus Howden 1976: 25. Colombia.
P. flexilis Champion 1911: 191. Guatemala, Honduras, and Mexico.
P. florivorus Howden 1998: 368. Venezuela.
P. giganteus Howden 1976: 94. Colombia.
P. gracilis Howden 1976: 48. Venezuela.
P. griseus (Voss 1954: 233). Peru.
P. hadromerooides (Kirsch 1868: 234). Colombia.
P. haruspex Howden 1986: 321. Mexico.
P. henryi Howden 1959: 398. *Mexico.
P. hercules Howden 1976: 109. Colombia and Venezuela.
P. hieroglyphicus Champion 1911: 190. Costa Rica.
P. hirtipes Champion 1911: 196. Mexico.
P. hispidus Champion 1911: 194. Mexico.
P. humboldti Howden 1976: 45. Colombia.
P. hystrix Howden 1976: 54. Venezuela.
P. inflatus Champion 1911: 197. Mexico.
P. julius Howden 1976: 27. Venezuela.
P. kirschi (Faust 1892: 2). Venezuela.
P. laticeps Champion 1911: 197. Mexico.
P. latirostris (Hustache 1938: 12). Argentina and Paraguay.
P. liberalis Howden 1976: 77. Venezuela.
P. longicollis Champion 1911: 206. *Costa Rica and Mexico.
P. longipenis Howden 1976: 32. Venezuela.
P. maculatus (Hustache 1923: 286). Argentina and Bolivia.
P. maculicollis Champion 1911: 200. Mexico.
P. mesosternalis Howden 1996: 895. Bolivia and Peru.
P. microcephalus Champion 1911: 205. Mexico.

- P. minax* Dohrn 1880: 157. Colombia.
- P. mirirostris* Howden 1976: 23. Colombia.
- P. modestus* (Faust 1892: 3). Colombia and Venezuela.
- P. multiseriatus* Howden 1976: 121. Venezuela.
- P. naupactoides* (Pascoe 1881: 38). Brazil (?) and Colombia.
- P. nodifer* Champion 1911: 206. *Colombia, El Salvador, Honduras, Jamaica, Nicaragua, and Venezuela.
- P. notabilis* Howden, 1976: 128. Colombia.
- P. novagranadæ* Howden 1976: 80. Colombia and Venezuela.
- P. obliquus* Champion 1911: 193. Mexico.
- P. olympus* Howden 1976: 57. Colombia.
- P. opalescens* (Faust 1892: 2). Panama and Venezuela.
- P. ornatifrons* Champion 1911: 198. Mexico.
- P. peckorum* Howden 1976: 125. Colombia and Venezuela.
- P. penai* Howden 1996: 914. Ecuador.
- P. peruvianus* Voss 1954: 230. Peru.
- P. pilosipctus* Howden 1976: 159. Colombia and Ecuador.
- P. platensis* Brèthes 1913: 192. Argentina, Brazil, and Paraguay.
- P. plumosiventris* Howden 1959: 401. *Mexico.
- P. pollinosus* Howden 1996: 892. Argentina, Bolivia, Brazil, Ecuador, and Peru.
- P. porosus* (Bohemian 1840: 294). Argentina, Brazil, and Paraguay.
- P. procollis* Howden 1976: 84. Colombia and Venezuela.
- P. pygmaeus* Howden 1976: 70. Colombia.
- P. quadrinodosus* Champion 1911: 199 [= *P. mexicanus* Voss 1954: 230, **syn. nov.**]. Guatemala and Mexico.
- P. quichensis* Champion 1911: 199. Guatemala.
- P. reductus* Howden 1976: 141. Colombia.
- P. regina* Howden 1976: 177. Colombia and Venezuela.
- P. robustulus* Emden 1939: 257 [replacement name for *P. robustus* (Schaeffer 1908: 215), not Faust 1892]. *Mexico (?).
- P. rotundicollis* (Fall 1907: 262) [= *P. deppressus* Pierce 1913: 403; *P. bryanti* Tanner 1907: 262]. *Mexico.
- P. santaanae* Howden 1976: 151. Venezuela.
- P. santamartae* Howden 1976: 165. Colombia.
- P. scorpioides* Howden 1996: 903. Peru.
- P. scutellatus* Howden 1976: 97. Colombia.
- P. semiconnatus* Howden 1996: 908. Argentina and Bolivia.
- P. separatus* Howden 1976: 51. Colombia.
- P. signatus* Voss 1954: 229. Brazil.
- P. sinuatipes* Champion 1911: 204. Mexico.
- P. smithsoni* Howden 1996: 919. Peru.
- P. spinipennis* Howden 1996: 900. Bolivia and Ecuador.
- P. sublineatus* Champion 1911: 203 [= *P. vitticollis* Champion 1911: 202, misident., in part]. Costa Rica, Grenada, Panama, and St. Vincent.
- P. submaculatus* Voss 1932: 34. Guatemala.
- P. subtilis* Howden 1996: 887. Colombia, Peru, and Venezuela.
- P. summus* Howden 1976: 143. Colombia.

- P. tessellatus* Howden 1976: 189. Colombia.
- P. testaceipes* Hustache 1929: 181. Dominican Republic, Grenada, Guadeloupe, and St. Vincent.
- P. thomasi* Howden 1996: 889. El Salvador, Guatemala, Honduras, and Mexico.
- P. tibialis* Boheman 1840: 297. Mexico.
- P. tinctorius* Howden 1976: 116. Colombia.
- P. torosus* Howden 1996: 910. Argentina, Brazil, and Paraguay.
- P. torquatus* Howden 1976: 62. Colombia.
- P. truncatus* Howden 1976: 138. Colombia.
- P. undatus* Champion 1911: 201. Guatemala.
- P. upsilon* Howden 1976: 90. Colombia.
- P. varicolor* Champion 1911: 193. Guatemala.
- P. variegatus* (Pierce 1915: 11). Peru.
- P. variogenitus* (Guenther 1933: 113) [= *P. variegatus* (Hustache 1926: 157), not Pierce 1915].
Argentina.
- P. viridiventris* Champion 1911: 199. Honduras and Mexico.
- P. vitticollis* Champion 1911: 202. Colombia, Guatemala, Honduras, Mexico, and Nicaragua.
- Paradacrys* Howden 1970: 40
- P. ensiformis* Howden 1970: 43. Bahamas.
- P. spatulatum* Howden 1970: 41. Bahamas, Turks, and Caicos Id.
- Paululusus* Howden 1970: 32
- P. calypso* Howden 1970: 34. Haiti.
- P. constanzae* Howden 1970: 38. Dominican Republic.
- P. hispaniolae* Howden 1970: 36. Dominican Republic and Haiti.
- Platyaspistes* Schoenherr 1840: 396 [= *Platyaspistes* Blanchard 1851: 319, lapsus; *Platyaspites* Perroud 1853: 492, lapsus]
- P. argentinensis* Kuschel 1950: 11 [= *P. alternans* Bruch 1915: 409, misident., not Fahraeus 1840]. Argentina.
- P. glaucus* Fahraeus 1840: 399. Chile.
- P. marginalis* Fahraeus 1840: 401 [= *P. validus* Chevrolat 1879: CXXVI]. Chile.
- P. prasinus* (Erichson 1834: 259) [= *P. limbatus* Chevrolat 1879: CXXVI]. Chile.
- P. unicolor* Perroud 1853: 492 [= *P. lateralis* Chevrolat 1879: CXXVI]. Chile.
- P. venustus* (Erichson 1834: 260). Chile.
- Scalaventer* Howden 1970: 3
- S. caymani* Howden 1970: 22. Grand Cayman.
- S. cocolobae* Howden 1970: 10. Jamaica.
- S. convexifrons* Howden 1970: 26. Bahamas.
- S. cubensis* Howden 1970: 31. Cuba.
- S. cyrillae* Howden 1970: 7. Jamaica.
- S. gelinasus* Howden 1970: 27. Dominican Republic.
- S. jamaicensis* Howden 1970: 24. Jamaica.
- S. litoreus* Howden 1970: 14. Jamaica.
- S. montanus* Howden 1970: 17. Jamaica.
- S. remotus* Howden 1970: 19. Jamaica.

S. subtropicus (Fall 1907: 263). *Cuba and Mexico.
S. valkyrius Howden 1970: 29. Jamaica.

Tanymecus Germar 1817: 341 [= *Tamymecus* Thomson 1859: 131, lapsus; *Tanycemus* Hamilton 1895: 344, lapsus; *Tanymoeclus* Brimley 1938: 235, lapsus]

T. cervinus Fahraeus 1840: 237. Brazil.

T. confusus Say 1831: 9 [= *T. confertus* Gyllenhal 1834: 88; *T. variabilis* Fahraeus 1840: 241].

*Belize, Guatemala, Honduras, Mexico, and Nicaragua.

T. hirsutus Champion 1911: 179. Mexico.

Subtribe NAUPACTINA Lacordaire 1863

Cyphina Lacordaire 1863.

Neocyphina Hustache 1919.

Platyomina Voss 1934.

Acyphus Heller 1921: 21

A. renggeri (Labram & Imhoff 1849: no. 61) [= *A. funicularis* Heller 1921: 27]. Argentina, Brazil, Paraguay, and Uruguay.

Amitrus Schoenherr 1840: 929 [= *Canephorus* Kirsch 1889, not Erichson 1846; *Canephorulana* Strand 1943: 96; *Canedorus* Marshall 1946: 93; *Canephorotomus* Voss 1954: 212; *Pernaupactus* Voss 1954: 216]

A. alutaceus (Erichson 1834: 256) [= *A. rufipes* (Kirsch 1889: 16); *A. proximus* Voss 1954: 215, **syn. nov.**]. Bolivia and Peru.

A. coriaceus Erichson 1847: 128. Peru.

A. jelskyi (Kirsch 1889: 16) [= *A. ielskyi* Hustache 1947: 107, lapsus; *A. corvinus* (Voss 1954: 213)]. Bolivia and Peru.

A. mundus (Erichson 1834: 257) [= *A. sobrinus* (Voss 1954: 213), **syn. nov.**]. Peru.

A. nitens Voss 1947: 45. Bolivia and Peru.

A. rugicollis Voss 1954: 215. Peru.

A. titschacki Voss 1954: 217. Peru.

Aptolemus Schoenherr 1843: 1

A. rufipes Boheman 1843: 2. Argentina, Brazil, and Paraguay.

Aramigus Horn 1876: 93 [= *Aomopactus* Horn 1876: 94; *Pseudeudius* Voss 1934: 72]

A. conirostris (Hustache 1947: 117). Argentina, Brazil, and Uruguay.

A. curtulus (Hustache 1947: 118). Argentina.

A. globoculus Lanteri 1994: 120. Brazil.

A. intermedius Lanteri 1994: 121. Brazil.

A. pilosus Lanteri 1994: 118. Brazil.

A. planioculus Lanteri 1994: 120. Argentina and Brazil.

A. tessellatus (Say 1824: 318) [= *A. durius* (Germar 1824: 417); *A. pallidus* Horn 1876: 94; *A. chevrolati* (Voss 1934: 64); *A. griseus* (Voss 1934: 63); *A. biseriatus* (Hustache 1947: 117); *A. viridipallens* (Hustache 1947: 137); *A. santafecinus* Lanteri 1987: 181]. Argentina, Brazil, and Uruguay. Introduced into Chile and the United States.

A. uruguensis Normark & Lanteri 1996 : 312. Uruguay.

Artipus Sahlberg 1823: 22 [= *Artipus* Schoenherr 1823: col. 1141, not Sahlberg 1823]

A. calceatus Marshall 1926: 530. Haiti.

A. coryaceus Sahlberg 1823: 22. St. Barthélemy.

A. floridanus Horn 1876: 92. *Bahamas.

A. freyanus Kuschel 1958: 795. Jamaica.

A. grisescens Chevrolat 1880: 253. Jamaica.

A. monae Wolcott 1941: 102. Mona and Puerto Rico.

A. porosicollis Chevrolat 1880: 254. Bahamas and Cuba.

A. psittacinus Gyllenhal 1834: 95. Hispaniola.

A. unguiculatus Chevrolat 1880: 254. Cuba.

Asymmathetes Wibmer & O'Brien 1986: 53 [= *Asymmathetes* Voss 1954: 210, not available, type species not fixed]

A. aequatorialis (Kirsch 1889: 13). Ecuador.

A. episternalis (Kirsch 1889: 14) [= *A. excisipennis* Voss 1954: 210]. Ecuador.

A. nigrans (Olliff 1891: 66). Ecuador.

A. pascoei (Kirsch 1889: 12). Ecuador.

A. rugicollis Voss 1954: 210. Ecuador.

A. steinheili (Kirsch 1889: 13). Colombia.

A. vulcanorum (Kirsch 1889: 17) [= *A. brevis* (Olliff 1891: 68); *A. pygmaeus* (Olliff 1891: 68)]. Colombia and Ecuador.

Asynonychus Crotch 1867: 388

A. cervinus (Boheman 1840: 17) [= *A. godmanni* Crotch 1867: 389; *A. fulleri* (Horn 1876: 94);

A. simplex (Pascoe 1881: 39); *A. olindae* (Perkins 1900: 130); *A. goldmani* Viana 1937: 107, lapsus]. Argentina, Brazil, Chile, Paraguay, and Uruguay. Introduced into Chile (Isla de Pascua), Central and North America, West Indies, and Old World.

Atrichonotus Buchanan 1939: 15 [= *Floresianellus* Hustache 1939: 41, not available, type species not fixed; *Floresianus* Hustache 1939: 39; *Floresianellus* Lanteri 1980: 266]

A. convexifrons (Hustache 1939: 41). Argentina.

A. marginatus (Hustache 1939: 42). Argentina and Uruguay.

A. obscurus (Hustache 1947: 121). Argentina and Uruguay.

A. pacificus Kuschel 1958: 788. Chile.

A. sordidus (Hustache 1939: 40). Argentina, Brazil, and Uruguay. Introduced into Australia and New Zealand.

A. taeniatus (Berg 1881: 61) [= *A. texanus* (Pierce 1911: 49); *A. pictipennis* (Hustache 1947: 122)]. Argentina, Chile, and Uruguay. Introduced into North America, Australia, and New Zealand.

A. whiteheadi Lanteri 1995: 108. Argentina.

Brachystyloides Hustache 1923: 288

B. meskendahli (Voss 1954: 208). Peru.

B. pilosus Hustache 1923: 288 [= *B. bolivianus* (Voss 1932: 36)]. Argentina and Bolivia.

B. rotundatus Kuschel 1945: 120. Argentina.

Chamaelops Kirsch 1868: 235 [= *Temnoscapus* Chevrolat 1878: LV]
C. fissirostris (Chevrolat 1878: LV). Colombia.

C. munitus Kirsch 1868: 236. Colombia.

Curiades Pascoe 1880: 420
C. boisduvali (Bohemian 1840: 163). Brazil.

Cyphopsis Roelofs 1879: LII
C. clathrata Roelofs 1879: LII [= *C. caldeirai* (Bondar 1949: 174)]. Brazil.
C. jekelii Roelofs 1879: LII. Brazil.

Cyrtomon Schoenherr 1823: col. 1140 [= *Cyphus* Germar 1824: 427; *Neocyphus* Bedel 1883: 23]
C. gibber (Pallas 1781: 32) [= *C. argillaceus* (Germar 1824: 429); *C. chevrolati* (Bohemian 1840: 142)]. Brazil.
C. glaucus (Bovie 1907: 326). Argentina.
C. hirsutus (Hustache 1939: 44). Argentina.
C. inhalatus (Germar 1824: 430) [= *C. intermedius* (Hustache 1947: 19)]. Argentina, Bolivia,
Brazil, Paraguay, and Uruguay.
C. luridus (Bohemian 1840: 144). Argentina, Brazil, and Paraguay.
C. ovalipennis (Hustache 1938: 95). Argentina.
C. pistor (Bohemian 1833: 621). Brazil.

Enoplopactus Heller 1921: 21 [= *Naupactosis* Heller 1921: 21; *Cyphodellus* Hustache 1939: 39]
E. brunneomaculatus Hustache 1926: 158. Argentina.
E. catamarcensis Lanteri 1990: 81. Argentina.
E. hylula (Heller 1921: 27). Argentina.
E. lizeri (Hustache 1926: 163). Argentina.
E. ortizi (Blanchard 1891: 493) [= *E. heterothorax* Heller 1921: 25]. Argentina and Paraguay.
E. sanjuaninus Lanteri 1990: 83. Argentina.
E. sulfureovittatus Heller 1921: 26. Argentina.

Ericydeus Pascoe 1880 [= *Cyphus* of authors, not Germar 1824]
E. argentinensis Lanteri 1995: 406. Argentina.
E. bahiensis Lanteri 1995: 401. Brazil.
E. cupreolus Lanteri 1995: 410. Mexico.
E. duodecimpunctatus (Champion 1911: 237). Mexico.
E. forneri (Champion 1911: 238). Mexico.
E. hancocki (Kirby 1819: 435). Brazil.
E. laetus (LeConte 1854: 85). *Mexico.
E. modestus (Gyllenhal 1833: 625). Mexico.
E. nigropunctatus (Chevrolat 1877: 170) [= *E. humeralis* Hustache 1938: 95]. Brazil, Colombia,
Ecuador, French Guiana, Peru, and Venezuela.
E. placidus (Horn 1876: 90). *Mexico.
E. quadripunctatus (Champion 1911: 238). Mexico.
E. roseiventris (Champion 1911: 239). Honduras and Mexico.
E. schoenherri (Perty 1832: 73) [= *E. latreillei* (Bohemian 1833: 624); *E. illustris* (Chevrolat 1833: 140)]. Argentina, Brazil, French Guiana, and Venezuela.

E. sedecimpunctatus (Linnaeus 1758: 386) [= *E. hilaris* (Perty 1832: 74)]. Argentina, Bolivia, Brazil, Colombia, French Guiana, Guyana, and Venezuela.
E. viridans (Bohemian 1840: 149). Mexico.
E. yucatanus (Champion 1911: 239). Mexico.

Eurymetopus Schoenherr 1840: 112 [= *Metoponeurus* Gemminger 1871: 2188; *Eumetopus* Bosq 1935: 330, lapsus]
E. birabeni Kuschel 1945: 127. Argentina and Uruguay. Introduced into Australia.
E. bucki Kuschel 1945: 129. Brazil.
E. fallax Boheman 1840: 113. Argentina, Brazil, and Uruguay.
E. globosus Lanteri 1984: 274. Argentina.
E. oblongus (Hustache 1938: 16). Argentina.
E. unicolor (Hustache 1938: 16). Uruguay.
E. vittatus Kuschel 1945: 128. Uruguay.

Galapaganus Lanteri 1992: 230
G. ashlocki Lanteri 1992: 245. Ecuador (Galapagos Ids.).
G. blairi (Van Dyke 1953: 146). Ecuador (Galapagos Ids.).
G. caroli (Van Dyke 1953: 147). Ecuador (Galapagos Ids.).
G. collaris Lanteri 1992: 246. Ecuador (Galapagos Ids.).
G. conwayensis (Mutchler 1938: 15). Ecuador (Galapagos Ids.).
G. crockeri (Van Dyke 1953: 147). Ecuador (Galapagos Ids.).
G. darwini Lanteri 1992: 236. Ecuador (Galapagos Ids.).
G. femoratus Lanteri 1992: 234. Ecuador.
G. galapagoensis (Linell 1898: 268). Ecuador (Galapagos Ids.).
G. howdenae Lanteri 1992: 235. Ecuador.
G. lacertosus (Erichson 1847: 128). Peru.
G. propinquus (Bohemian 1859: 161). Ecuador.
G. squamosus Lanteri 1991: 247. Peru.
G. vandykei Lanteri 1992: 237. Ecuador (Galapagos Ids.).
G. williamsi (Van Dyke 1953: 150). Ecuador (Galapagos Ids.).

Hadropus Schoenherr 1826: 111
H. albiceris (Germar 1824: 432) [= *H. nubiculosus* Schoenherr 1826: 111; *H. albinus* Rosenschoeld 1840: 154; *H. fascicularis* Hustache 1938: 94]. Brazil and Paraguay.

Hoplopactus Chevrolat 1879: LXXXIV [= *Diaphorus* Faust 1892: 7; *Diapherontus* Berg 1898: 17; *Hoplopactus* Heller 1921: 21, lapsus; *Anaphorus* Marshall 1935: 515; *Diaperontus* Blackwelder 1947: 793, lapsus]
H. albarius (Faust 1892: 6). Venezuela.
H. armatus (Kirsch 1874: 385). Bolivia and Peru.
H. dentipes (Kirsch 1868: 231). Colombia.
H. griseus (Kirsch 1874: 388). Peru.
H. injucundus (Bohemian 1833: 601) [= *H. temperans* (Bohemian 1833: 610)]. Brazil and French Guiana.
H. lateralis Arrow 1903: 251. Bolivia and Brazil.
H. ochreatus Hustache 1947: 28. Bolivia.

H. pavidus (Boheman 1840: 107) [= *H. inustus* Chevrolat 1879: LXXXV; *H. morosus* (Faust, 1892: 7); *H. defolians* Marshall 1948: 97]. Colombia, Trinidad, and Venezuela.

H. rufipes (Kirsch 1868: 230). Colombia.

H. segnipes (Olliff 1891: 65). Ecuador.

H. signatus (Boheman 1833: 599). Brazil.

H. singularis Hustache 1938: 276. Bolivia and Peru.

H. suturalis (Kirsch 1868: 230). Colombia.

H. virens Marshall 1948: 97. Trinidad.

Lamprocyphopsis Lanteri 1990: 416

L. paraguayensis Lanteri 1990: 420. Paraguay.

L. viridimilensis (Kuschel 1950: 546). Argentina and Paraguay.

Lamprocyphus Marshall 1922: 184

L. augustus (Illiger 1802: 174) [= *L. angustus* Lacordaire 1863: 115, lapsus]. Brazil.

L. elegans (Roelofs 1879: LII). Brazil.

L. germari (Boheman 1833: 627). Brazil.

L. margaritaceus (Sturm 1826: 75) [= *L. gloriandus* (Schoenherr 1826: 108); *L. spixii* (Perty 1832: 74); *L. consularis* (Chevrolat 1838: 56)]. Brazil.

L. oliveirae (Roelofs 1879: LII). Brazil.

L. varnhageni (Germar 1824: 437) [= *L. jacobi* Hustache 1947: 18]. Argentina, Brazil, and Paraguay.

Litostylodes Hustache 1947: 14

L. chlorostomus (Germar 1824: 430). Brazil.

Litostylus Faust 1894: 368 [= *Leptostylus* Faust 1892: 8; *Germariella* Champion 1911: 234]

L. bovelli (Marshall 1922: 184). Barbados and Dominican Republic.

L. diadema (Fabricius 1787: 116) [= *L. juvencus* (Olivier 1807: 352); *L. peruvianus* Hustache 1947: 33]. Argentina, Brazil, Colombia, French Guiana, Honduras, Nicaragua, Panama, Peru, and Venezuela.

L. leucocephalus (Chevrolat 1880: 213). Guadeloupe.

L. marginicollis (Chevrolat 1880: 197) [= *L. viridis* (Chevrolat 1880: 197)]. Martinique.

L. pudens (Boheman 1833: 623). Antigua, Montserrat, St. Barthélemy, and St. Vincent.

L. strangulatus (Chevrolat 1880: 213) [= *L. triangulatus* Guenther & Zumpt 1933: 116; *L. triangularis* Blackwelder 1947: 793, lapsus]. Dominican Republic, Guadeloupe, and Montserrat.

L. suturalis (Faust 1892: 8). Venezuela.

Macrostylus Boheman 1840: 921 [= *Amphideritus* Schoenherr 1840: 117; *Mimographus* Schoenherr 1847: 34; *Pterotropis* Schoenherr 1847: 28; *Mimographopsis* Champion 1911: 229; *Steirarhinus* Champion 1911: 222; *Pantomorophilus* Voss 1954: 220]

M. albiventris Bordón 1991: 26. Venezuela.

M. alecsandri Bordón 1991: 18. Venezuela.

M. amandus (Kirsch 1868: 225) [= *M. argutulus* (Kirsch 1868: 228); *M. jekelii* (Kirsch 1868: 226); *M. lugens* (Kirsch 1868: 227); *M. moestus* (Kirsch 1868: 227); *M. viridanus* (Kirsch 1868: 226); *M. nudimaculatus* Voss 1953: 60]. Colombia and Panama.

- M. anabelleae* Bordón 1991: 16. Venezuela.
M. andreinae Bordón 1991: 20. Venezuela.
M. ardosiacus (Kirsch 1968: 229). Colombia.
M. aroensis Bordón 1991: 23. Venezuela.
M. avilensis Bordón 1991: 8. Venezuela.
M. barinensis Bordón 1991: 22. Venezuela.
M. beatricis Bordón 1991: 21. Venezuela.
M. bicinctus (Champion 1911: 227). Panama.
M. boconoensis Bordón 1991: 19. Venezuela.
M. brachyderoides (Hustache 1938: 275). Peru.
M. brevisetosus (Kirsch 1889: 15). Ecuador.
M. chilensis Kuschel 1949: 14. Chile.
M. cinctipennis (Champion 1911: 228). Guatemala.
M. cinereoguttatus (Champion 1911: 228). Colombia, Venezuela, and Central America.
M. coloniae Bordón 1991: 11. Venezuela.
M. crinitus Boheman 1840: 923. Brazil.
M. decolor (Boheman 1840: 106) [= *S. unicolor* (Chevrolat 1879: LXXVII); *S. conicollis* (Champion 1911: 223)]. Colombia, Guatemala (?), and Mexico.
M. depresicollis (Boheman 1840: 66). Colombia.
M. dietzi Bordón 1991: 23. Venezuela.
M. elegantulus (Boheman 1840: 64). Colombia.
M. falconensis Bordón 1991: 24. Venezuela.
M. guaniamensis Bordón 1991: 12. Venezuela.
M. guatimaltecus (Champion 1911: 226). Guatemala.
M. guatopensis Bordón 1991: 10. Venezuela.
M. hirtus (Voss 1953: 60). Colombia.
M. infucatus (Champion 1911: 224). Belize.
M. lacunitae Bordón 1991: 12. Venezuela.
M. laesicollis (Schoenherr 1847: 35). Colombia and Venezuela.
M. leporinus (Erichson 1847: 128) [= *M. peruvianus* (Brèthes 1920: 39)]. Bolivia and Peru.
M. lineatocollis (Champion 1911: 227) [= *M. cupreotinctus* (Champion 1911: 226); *M. filicornis* (Voss 1954: 225)]. Costa Rica, Panama, and Peru.
M. litoralis Bordón 1991: 27. Venezuela.
M. loscaracasensis Bordón 1991: 9. Venezuela.
M. mexicanus (Champion 1911: 231). Mexico.
M. micaceus (Hustache 1947: 26) [= *M. villosipennis* (Hustache 1947: 27)]. Argentina, Brazil, and Paraguay.
M. micans (Kirsch 1868: 229). Colombia.
M. mucuyensis Bordón 1991: 14. Venezuela.
M. nebulosus (Champion 1911: 224). Costa Rica and Panama.
M. nocens (Kuschel 1958: 135). Costa Rica and Nicaragua.
M. ocellatus Lanteri 1985: 4. Argentina and Brazil.
M. pauper (Olliff 1891: 66). Ecuador.
M. peninsularis Bordón 1991: 14. Venezuela.
M. pittieri Bordón 1991: 6. Venezuela.
M. plicaticollis (Hustache 1938: 274) [= *M. ochraceus* (Hustache 1938: 274), **syn. nov.**]. Colombia.

- M. puberulus* (Boheman 1859: 120). Chile and Peru.
M. pustulatus (Champion 1911: 230). Costa Rica and Panama.
M. rotundicollis (Kirsch 1889: 15). Colombia.
M. rugicollis (Kirsch 1868: 223). Colombia.
M. santanae Bordón 1991: 26. Venezuela.
M. serripes (Champion 1911: 225). Costa Rica and Panama.
M. setosus (Schoenherr 1847: 28) [= *M. squamosus* Kirsch 1868: 224]. Colombia.
M. tenuicornis (Champion 1911: 226). Panama.
M. tomentosus (Boheman 1833: 545) [= *M. cinereus* (Boheman 1840: 886); *M. peruvianus* (Boheman 1840: 885); *M. peruanus* (Erichson 1847: 127)]. Peru.
M. trujillanus Bordón 1991: 19. Venezuela.
M. uliani Bordón 1991: 25. Venezuela.
M. valeranus Bordón 1991: 16. Venezuela.
M. varians (Boheman 1840: 65). Colombia.
M. versicolor Kirsch 1889: 15. Colombia.
M. vilis (Boheman 1840: 118) [= *M. vittis* Heller 1921: 20, lapsus]. Colombia.
M. viridicans (Champion 1911: 229). Costa Rica and Panama.
M. viridiventris Bordón 1991: 17. Venezuela.
M. vittatus (Kirsch 1889: 14). Colombia.
M. yacambuanus Bordón 1991: 5. Venezuela.

Megalostyloides Champion 1911: 246

M. hirsutus Champion 1911: 247. Honduras, Mexico, and Nicaragua.

Megalostylus Schoenherr 1840: 114 [= *Hadrotomus* Dejean 1834: 253, nom. nud.]

M. albicans Lacordaire, unknown date: pl. 62 [= *M. farinosus* Chevrolat 1878: LCVI, syn. nov.; *M. expansus* Pascoe 1881: 42, syn. nov.]. Mexico.

M. brevipilis Champion 1911: 243. Mexico.

M. dilaticollis Champion 1911: 246. Mexico.

M. fusiformis Champion 1911: 245. Mexico.

M. macrophthalmus Champion 1911: 246. Mexico.

M. rhodopus Boheman 1840: 116. Mexico.

M. splendidus Chevrolat 1878: LV [= *M. physops* Champion 1911: 245, syn. nov.]. Mexico.

M. sturmi Boheman 1840: 115 [= *M. villosus* Chevrolat 1878: LIV, syn. nov.]. Mexico.

M. tomentosus Champion 1911: 243. Mexico.

Melanocyphus Jekel 1875: 143 [= *Melactus* Pascoe 1886: 416; *Melanocythus* Hustache 1947: 15, lapsus]

M. bispinus (Boheman 1840: 150). Colombia.

M. lugubris (Boheman 1840: 147). Colombia.

Mendozella Hustache 1939: 42

M. curvispinnis Hustache 1926: 161. Argentina.

Miocyphus Hustache 1939: 38 [= *Myocyphus* Hustache 1939: 38, lapsus]

M. laticeps Hustache 1939: 38. Brazil and French Guiana.

Mionarthrus Kuschel 1950: 11

M. cinerascens (Blanchard 1851: 318). Chile.

Moropactus Marshall 1938: 1

M. leucospilus (Germar 1824: 422) [= *M. lepidotus* Marshall 1938: 3]. Brazil.

M. perfidus Marshall 1938: 3. Brazil.

Naupactus Dejean 1821: 94 [= *Alceis* Billberg 1820: 45; *Iphius* Schoenherr 1823: col. 1140; *Leptocerus* Germar 1824: 417; *Pantoplanes* Schoenherr 1840: 111; *Antelmia* Hustache 1919: 464; *Archopactus* Heller 1921: 20; *Graphognathus* Buchanan 1939: 11; *Archiopactus* Monrós 1944: 181, lapsus; *Naupactuosus* Hustache 1947: 108, lapsus]

N. acerbus Boheman 1840: 31. Brazil.

N. aerosus Boheman 1840: 2. Brazil.

N. agglomeratus Hustache 1938: 269. Bolivia.

N. albidus Perroud 1853: 461. Brazil.

N. albolateralis Hustache 1947: 78. Argentina.

N. albopunctatus Boheman 1840: 53. Brazil.

N. albovittatus Voss 1932: 34. Bolivia.

N. albulus Boheman 1833: 609. Brazil, French Guiana, and Suriname.

N. alternevittatus Hustache 1947: 95. Uruguay.

N. ambiguus Boheman 1840: 16. Brazil and Paraguay.

N. ambitiosus Boheman 1833: 603. Brazil.

N. amens Boheman 1833: 576. Brazil.

N. anceps Boheman 1833: 574. Brazil.

N. ancora Marshall 1949: 321. Brazil.

N. angulithorax Hustache 1947: 64. Argentina.

N. annae Bordón 1997: 35. Venezuela.

N. anthribiformis Boheman 1833: 601. Brazil.

N. aptus Boheman 1840: 5 [= *N. aurulans* Boheman 1849: 82; *N. bistigma* Boheman 1840: 84; *N. roseiventris* Boheman 1840: 87; *N. scintillans* Boheman 1840: 80; *N. tibialis* Boheman 1840: 97]. Brazil and Paraguay.

N. argentatus Hustache 1947: 91. Argentina.

N. argentinensis (Hustache 1926: 91) [= *N. brevipubens* (Hustache 1926: 160)]. Argentina.

N. argyrostomus Boheman 1840: 105. Brazil.

N. aulacus (Germar 1824: 421). Brazil.

N. aurichalceus Boheman 1833: 603 [= *N. amoenus* Boheman 1833: 604; *N. mitis* Boheman 1833: 606; *N. psittacinus* Boheman 1840: 90; *N. ruricola* Boheman 1833: 608; *N. suavis* Boheman 1833: 605]. Brazil.

N. auricinctus Boheman 1833: 584 [= *N. chloris* Hustache 1947: 82]. Argentina, Brazil, and Paraguay.

N. auripes (Hustache 1947: 125). Argentina.

N. aurolimbatus Boheman 1840: 41 [= *N. auromarginatus* Boheman 1840: 42]. Brazil.

N. balteus Voss 1934: 67. Brazil.

N. barbicauda Boheman 1840: 79. Brazil.

N. basalis Hustache 1947: 81. Bolivia.

N. basilicus (Germar 1824: 421) [= *N. auriventris* (Germar 1824: 425); *N. granicollis* Boheman 1833: 597; *N. balteatus* Boheman 1840: 77; *N. quadrigatus* Boheman 1840: 78; *N. viridiustulatus* Boheman 1840: 86]. Brazil.

- N. bellus* Boheman 1833: 598. Brazil.
- N. bipes* (Germar 1824: 426). Bolivia, Brazil, and Paraguay.
- N. bipunctatus* Boheman 1833: 607 [= *N. binotatus* Boheman 1840: 100; *N. calvus* Boheman 1840: 100]. Brazil and Paraguay.
- N. bohumilae* Bordón 1997: 29. Venezuela.
- N. bondari* Marshall 1937: 469. Brazil.
- N. bosqi* Hustache 1947: 80. Argentina and Brazil.
- N. brevicrinitus* Hustache 1947: 103. Argentina.
- N. bridgesii* Waterhouse 1844: 54. Argentina and Bolivia.
- N. bruchi* (Heller 1921: 23) [= *N. niveopectus* (Hustache 1926: 160)]. Argentina.
- N. brunneus* (Hustache 1947: 123). Argentina and Bolivia.
- N. calamuchitanensis* Hustache 1947: 65. Argentina.
- N. camachoi* Bordón 1997: 33. Venezuela.
- N. carinirostris* (Hustache 1947: 134). Argentina.
- N. castaneus* Hustache 1947: 90. Argentina and Bolivia.
- N. celator* Boheman 1833: 607. Brazil and French Guiana.
- N. cephalotes* (Hustache 1947: 144). Argentina and Bolivia.
- N. chaconi* Bordón 1997: 19. Venezuela.
- N. chalybeipes* Boheman 1840: 47. Brazil.
- N. chedasi* Bordón 1997: 21. Venezuela.
- N. chevrolati* Boheman 1833: 85. French Guiana.
- N. chloraspis* Erichson 1847: 128. Peru.
- N. chloropleurus* Pascoe 1881: 39. Brazil.
- N. chordinus* Boheman 1833: 573 [= *N. suffitus* Boheman 1833: 573; *N. fernandezi* Hustache 1947: 99; *N. sericellus* Hustache 1947: 101; *N. sericeus* Hustache 1947: 100]. Argentina and Uruguay.
- N. cinerascens* (Perroud 1853: 454). Guatemala.
- N. cinereidorsum* Hustache 1947: 92. Argentina and Uruguay.
- N. cinerosus* Boheman 1833: 577 [= *N. marginalis* Boheman 1833: 577; *N. parsevali* (Costa-Lima 1936: 23); *N. inimicus* (Marshall 1938: 211); *N. roseomicans* (Hustache 1947: 129); *N. subbimaculatus* (Hustache 1947: 129); *N. ursius* (Hustache 1947: 130)]. Argentina, Brazil, and Uruguay.
- N. clavijoi* Bordón 1997: 36. Venezuela.
- N. concinnus* Boheman 1840: 48. Brazil.
- N. condecoratus* Boheman 1840: 38. Argentina, Brazil, and Paraguay.
- N. cupreus* Bordón 1997: 27. Venezuela.
- N. curialis* (Germar 1824: 423) [= *N. stultus* Boheman 1840: 73]. Brazil and French Guiana.
- N. curtus* Boheman 1833: 570 [= *N. unicolor* Boheman 1833: 570]. Brazil.
- N. curvilineus* Boheman 1840: 63. Brazil.
- N. cyphoides* (Heller 1921: 24) [= *N. griseomaculatus* (Hustache 1923: 280); *N. viridimarginalis* Hustache 1947: 43]. Argentina, Bolivia, and Uruguay.
- N. dapsilis* (Perty 1832: 73) [= *N. wintheimii* Boheman 1833: 595]. Brazil.
- N. decorus* (Fabricius 1775: 146). Brazil.
- N. delicatulus* Hustache 1947: 67. Argentina and Brazil.
- N. denudatus* Hustache 1923: 283 [= *N. vicinus* Hustache 1923: 284]. Argentina and Bolivia.
- N. deplorabundus* Boheman 1833: 609. Brazil.
- N. deses* Schoenherr 1840: 49. Brazil.

- N. dissimilis* Hustache 1947: 69. Argentina, Brazil, and Paraguay.
- N. dissimulator* Boheman 1840: 8 [= *N. fallax* Boheman 1840: 9]. Argentina, Brazil, Paraguay, and Uruguay.
- N. dives* Klug 1829: 6 [= *N. klugii* Boheman 1833: 572]. Argentina, Brazil, and Uruguay.
- N. elegans* Lucas 1857: 152. Brazil.
- N. faldermanni* Boheman 1840: 70 [= *N. ornatus* Boheman 1840: 88]. Brazil.
- N. fatuus* Boheman 1833: 588 [= *N. sulphureosignatus* Blanchard 1847: 202; *N. sulfureosignatus* Bosq 1935: 332, lapsus]. Bolivia and Brazil.
- N. fimbriatus* Hustache 1938: 271. Bolivia and Peru.
- N. fulgerens* Lucas 1857: 151 [= *N. fulgurans* Gemminger & Harold 1871: 2199, lapsus]. Brazil.
- N. fulvus* (Hustache 1947: 128). Argentina.
- N. fuscus* Boheman 1840: 54. Brazil.
- N. glaucus* (Perty 1832: 72). Brazil.
- N. habenatus* Marshall 1949: 322. Brazil.
- N. hirsuticeps* (Hustache 1947: 131). Argentina.
- N. hirsutus* Hustache 1947: 72. Argentina.
- N. hirtellus* (Voss 1932: 33) [= *N. cupreata* (Voss 1932: 33); *N. caroli* Hustache 1947: 70]. Argentina and Paraguay.
- N. humilis* (Hustache 1947: 119). Argentina.
- N. hypocrita* (Germar 1824: 420) [= *N. civica* Voss 1934: 64; *N. viridula* Voss 1934: 65]. Brazil.
- N. illotus* (Germar 1824: 424). Brazil.
- N. imbutus* Pascoe 1881: 40. Ecuador.
- N. impurus* Boheman 1833: 580. Brazil.
- N. inermis* Hustache 1947: 86 [= *N. paraguayanus* Hustache 1947: 87]. Argentina, Brazil, and Paraguay.
- N. insignis* Boheman 1840: 95. Unknown.
- N. instabilis* Boheman 1840: 71. Colombia and Venezuela.
- N. institor* Boheman 1833: 597 [= *N. rufipes* Boheman 1833: 589]. Brazil.
- N. jacobi* Hustache 1947: 85. Brazil and Paraguay.
- N. jekeli* Kirsch 1874: 386. Peru.
- N. jolyi* Bordón 1997: 33. Venezuela.
- N. lax* (Germar 1824: 427). Brazil.
- N. laticeps* (Champion 1911: 233). Costa Rica and Honduras.
- N. laticollis* Hustache 1947: 74. Argentina.
- N. latifrons* Boheman 1833: 592 [= *N. turmalinus* Boheman 1840: 59]. Brazil.
- N. lattkei* Bordón 1997: 37. Venezuela.
- N. leucogaster* (Perty 1832: 72). Bolivia, Brazil, and French Guiana.
- N. leucographus* Boheman 1840: 45. Brazil.
- N. leucoloma* Boheman 1840: 62 [= *N. dubius* (Buchanan 1942: 109); *N. pilosus* (Buchanan 1942: 107); *N. striatus* (Buchanan 1942: 108); *N. fecundus* (Buchanan 1947: 21); *N. imitator* (Buchanan 1947: 21)]. Argentina, Brazil, and Uruguay. Introduced into Australia, Chile (incl. Isla de Pascua), New Zealand, Peru, South Africa, and the United States.
- N. leucophaeus* Boheman 1833: 600. Brazil.
- N. lineatus* Boheman 1840: 3. Brazil.
- N. litoris* Bordón 1997: 30. Venezuela.
- N. lizeri* Hustache 1923: 280 [= *N. unicolor* Hustache 1923: 282, **syn. nov.**; *N. uniformis* Hustache 1947: 45, lapsus]. Bolivia.

- N. llanensis* Bordón 1997: 31. Venezuela.
- N. loripes* (Germar 1824: 419). Brazil.
- N. luteipes* (Hustache 1947: 130). Argentina and Paraguay.
- N. macilentus* Boheman 1833: 594 [= *N. lateralis* Boheman, 1840: 43]. Brazil.
- N. mariaeolosiae* Bordón 1997: 29. Venezuela.
- N. maritimus* Bordón 1997: 23. Venezuela.
- N. martinezii* Bordón 1997: 21. Venezuela.
- N. mimicus* Hustache 1938: 268 [= *N. justus* Bondar 1949: 176]. Brazil.
- N. minor* (Buchanan 1942: 107) [= *N. breviscapus* Hustache 1947: 102]. Argentina and Uruguay. Introduced into the United States.
- N. minutellus* (Wibmer & O'Brien 1986: 65) [= *N. minutus* (Voss 1954: 221), not Hustache 1938]. Peru.
- N. minutus* (Hustache 1938: 277). Bolivia.
- N. morio* Boheman 1833: 592. Brazil.
- N. mulsanti* Perroud 1853: 457. Brazil.
- N. navicularis* Boheman 1840: 23. Argentina and Brazil.
- N. navus* Marshall 1949: 324. Brazil.
- N. nubilosus* Boheman 1840: 51. Brazil.
- N. obsoletus* Kirsch 1874: 387. Bolivia and Peru.
- N. ocami* Bordón 1997: 28. Venezuela.
- N. ochreonotatus* Voss 1934: 66. Brazil.
- N. optatus* (Herbst 1797: 40) [= *N. ypsilon* Olivier 1807: 330; *N. ypsilon* Olivier 1807: 330, lapsus; *N. faustus* Erichson 1848: 568]. Brazil, Guyana, and Suriname.
- N. ornatus* (Pascoe 1879: 185) [= *N. longimanus* (Fabricius 1775: 147), not Forster 1771]. Brazil. Introduced into Ascencion Id.
- N. ovatus* (Hustache 1947: 124). Brazil.
- N. pedestris* Voss 1934: 65. Brazil.
- N. penai* Bordón 1997: 22. Venezuela.
- N. peregrinus* (Buchanan 1939: 14). Argentina. Introduced into Azores and the United States.
- N. peruvianus* Hustache 1938: 272. Bolivia and Peru.
- N. pictus* Boheman 1840: 24. Brazil and Paraguay.
- N. pilipes* Hustache 1947: 96. Uruguay.
- N. pithecius* (Germar 1824: 426) [= *N. beta* Schoenherr 1833: 584, not Germar 1824]. Bolivia and Brazil.
- N. plagiatus* Lucas 1857: 154. Brazil.
- N. polliger* Boheman 1833: 588 [= *N. flavipennis* Boheman 1840: 55]. Bolivia and Brazil.
- N. postfasciatus* (Hustache 1947: 141). Argentina, Paraguay, and Uruguay.
- N. postsignatus* (Voss 1940: 13). Brazil.
- N. praedatus* Erichson 1847: 128. Peru.
- N. prasinus* (Hustache 1947: 126), **comb. nov.** Argentina.
- N. proximus* Voss 1934: 65. Brazil.
- N. pulchellus* Kuschel 1958: 784. Brazil.
- N. pupulus* Boheman 1840: 81 [= *N. pulverulentus* Boheman 1840: 98]. Brazil.
- N. purpureoviolaceus* Hustache 1947: 98 [= *N. persimilis* (Hustache 1947: 140)]. Argentina, Paraguay, and Uruguay.
- N. rivulosus* (Olivier 1790: 545). Argentina, Brazil, and Paraguay.
- N. roborosus* (Germar 1824: 423). Brazil.

- N. romeroi* Bordón 1997: 25. Venezuela.
N. rosalesi Bordón 1997: 20. Venezuela.
N. roscidus Erichson 1848: 568. Guyana.
N. rubiginosus (Fabricius 1801: 521). Brazil, French Guiana, and Suriname.
N. ruficornis Boheman 1840: 40. Brazil.
N. rugosus Hustache 1947: 93. Argentina.
N. ruizi (Brèthes 1925: 204) [= *N. subvittatus* Fairmaire & Germain 1861: 7, not Boheman 1840; *N. subvittulus* Emden 1936: 23; *N. intermedius* (Hustache 1947: 140); *N. variabilis* (Hustache 1947: 139)]. Argentina and Chile.
N. sahlbergi Boheman 1840: 32. Brazil.
N. sanarensis Bordón 1997: 26. Venezuela.
N. sanfilippii Bordón 1997: 22. Venezuela.
N. sanguinipes Hustache 1938: 269. Brazil.
N. santanae Bordón 1997: 36. Venezuela.
N. scelestus Boheman 1833: 579. Brazil.
N. schapleri Hustache 1947: 76. Argentina and Paraguay.
N. schnusei (Voss 1954: 223). Peru.
N. sellatus Boheman 1840: 96 [= *N. obliquus* Boheman 1840: 96; *N. subvittatus* Boheman 1840: 102]. Brazil and Paraguay.
N. serenus Pascoe 1881: 40 [= *N. parallelus* Hustache 1947: 77]. Brazil and Paraguay.
N. setarius Boheman 1833: 578. Brazil.
N. sexmaculatus Lucas 1857: 153. Peru.
N. signatus (Blanchard 1847: 201). Bolivia.
N. signipennis Boheman 1840: 92 [= *N. enitescus* Boheman 1840: 99; *N. squameus* Boheman 1840: 94]. Brazil.
N. similis (Hustache 1947: 127). Argentina.
N. simulator (Voss 1954: 222). Peru.
N. sommeri Boheman 1840: 76. Brazil.
N. sparsus Boheman 1840: 103. Brazil and Paraguay.
N. stauropterus (Germar 1824: 419). Brazil.
N. stigmaticus Boheman 1840: 57 [= *N. argenteus* Hustache 1947: 89; *N. argentus* Hustache 1947: 57, lapsus]. Brazil.
N. subacutus Boheman 1840: 33. Brazil.
N. submaculatus Hustache 1947: 71. Brazil and Paraguay.
N. sulcipes Hustache 1947: 94. Uruguay.
N. sulfuratus (Champion 1911: 232). Mexico.
N. sulphureoviridis (Hustache 1938: 276). Argentina, Brazil, and Paraguay.
N. sulphurifer Pascoe 1881: 41 [= *N. interruptus* Hustache 1947: 42, **syn. nov.**]. Argentina, Paraguay, and Uruguay.
N. suturalis Boheman 1840: 52. Brazil.
N. tachirensis Bordón 1997: 20. Venezuela.
N. tarsalis Boheman 1840: 58 [= *N. glaucivittatus* Blanchard 1847: 202; *N. albividiventris* Hustache 1947: 75]. Argentina, Bolivia, Brazil, and Paraguay.
N. transversus Boheman 1840: 11. Argentina and Brazil.
N. tremolerasi Hustache 1947: 97. Brazil and Uruguay.
N. tucumanensis Hustache 1947: 79. Argentina, Bolivia, and Paraguay.
N. umbrinus Hustache 1947: 67. Paraguay.

- N. univittatus* Boheman 1833: 587. Brazil.
N. variegatus Hustache 1938: 267. Brazil.
N. variesignatus Voss 1951: 365. Brazil.
N. velox (Fabricius 1787: 117) [= *N. dasypus* Boheman 1840: 72]. French Guiana.
N. venezolanus Hustache 1938: 270. Colombia, Venezuela, and Central America.
N. verecundus Hustache 1947: 105 [= *N. calamuchitanus* Hustache 1947: 106; *N. vianai* Hustache 1947: 106]. Argentina and Chile.
N. versatilis Hustache 1947: 88 [= *N. imbellis* Hustache 1947: 89]. Argentina, Brazil, and Paraguay.
N. viduatus Boheman 1840: 60. South America.
N. vilmae Bordón 1997: 34. Venezuela.
N. viloriai Bordón 1997: 18. Venezuela.
N. virens Boheman 1840: 6. Argentina and Brazil.
N. virescens (Champion 1911: 233). Mexico.
N. viridicinctus Boheman 1833: 596. Argentina and Brazil.
N. viridicyaneus Hustache 1947: 84. Bolivia.
N. viridimicans Hustache 1923: 282. Bolivia and Peru.
N. viridinutens Hustache 1947: 66. Argentina.
N. viridiplagiatus Boheman 1840: 75. Brazil.
N. viridisquamosus Boheman 1840: 37. Brazil and Paraguay.
N. viridissimus (Hustache 1919: 465) [= *N. viridisquamosus* (Boheman 1859: 121), not Boheman 1840]. Argentina and Uruguay. Introduced into Mauritius.
N. viridulus Hustache 1947: 66. Argentina.
N. vittatus Mannerheim 1833: 591. Brazil.
N. vossi Morrone 1997: 12 [= *N. bondari* (Voss 1951: 366), not Marshall 1937]. Brazil.
N. wilsoni Boheman 1840: 50. Brazil.
N. xanthographus (Germar 1824: 424). Argentina, Brazil, Paraguay, and Uruguay. Introduced into Chile.

- Neoericydeus* Hustache 1938: 93 [= *Neoerycideus* Hustache 1938: 94, lapsus]
N. coelestinus (Rosenschoeld 1840: 203). French Guiana.
N. gratiosus Hustache 1938: 93. Bolivia, Brazil, and Ecuador.
N. tuberculatus (Hustache 1938: 73). Brazil.

- Pantomorus* Schoenherr 1840: 942 [= *Pantopactus* Horn 1876: 96; *Pantomerus* Chevrolat 1879: CXXX, lapsus; *Athetetes* Pascoe 1886: 415; *Athetes* Dalla Torre, Emden, & Emden 1936: 25, lapsus; *Pantamorus* Cortés 1941: 61, lapsus; *Pantomoris* Van Dyke 1953: 150, lapsus].
P. albicans Sharp 1891: 164. Mexico.
P. albosignatus Boheman 1840: 943. El Salvador, Guatemala, and Mexico.
P. annectens Sharp 1891: 163. Mexico.
P. asperatus Sharp 1891: 156. Mexico.
P. brevipes Sharp 1891: 163. Mexico.
P. circumcinctus Sharp 1891: 160. Guatemala and Honduras.
P. comes Kuschel 1956: 335. El Salvador.
P. crinitus (Boheman 1840: 20) [= *P. affinis* Sharp 1891: 157, **syn. nov.**]. El Salvador, Guatemala, and Mexico.
P. distans Sharp 1891: 162. Costa Rica and Panama.

- P. dorsalis* Sharp 1891: 159. Guatemala and Honduras.
P. faber Sharp 1891: 159. Costa Rica and Honduras.
P. facialis Sharp 1891: 156. Mexico.
P. femoratus Sharp 1891: 165 [= *P. robustus* Sharp 1891: 166]. Costa Rica, El Salvador, Honduras, and Nicaragua.
P. globicollis (Pascoe 1886: 416). Costa Rica, Guatemala, and Mexico.
P. horridus Champion 1911: 334. Mexico.
P. longulus Sharp 1891: 154. Mexico.
P. maculosus (Boheman 1840: 21). Mexico.
P. mollis Sharp 1891: 155. Mexico.
P. nobilis (Boheman 1840: 19). Mexico.
P. parvulus Sharp 1891: 153. Mexico.
P. picipes Sharp 1891: 154. Mexico.
P. picturatus Sharp 1891: 160. Costa Rica, Guatemala, Honduras, and Mexico.
P. rufidis Sharp 1891: 162. Costa Rica, Guatemala, Honduras, and Mexico.
P. rufipes Sharp 1891: 161. Mexico.
P. salvadorensis Kuschel 1956: 339. El Salvador.
P. salvini Sharp 1891: 155. Guatemala.
P. sobrinus Sharp 1891: 158. Guatemala.
P. strabo Sharp 1891: 165. Honduras and Nicaragua.
P. stupidus (Boheman 1840: 18). Guatemala and Mexico.
P. subcinctus Sharp 1891: 158. Guatemala.
P. sulfureus Champion 1911: 338. Mexico.
P. trituberculatus Champion 1911: 337. Mexico.
P. uniformis Sharp 1891: 161. Mexico.
P. viridicans Sharp 1891: 164. Mexico.
- Parapantomorus* Emden 1936: 25 [= *Pseudopantomorus* Heller 1921: 20]
P. fluctuosus (Boheman 1840: 15) [= *P. sharpi* (Heller 1921: 22)]. Argentina, Bolivia, Brazil, and Paraguay.
P. quatuordecimpunctatus (Hustache 1923: 284). Argentina, Bolivia, and Paraguay.
- Parasynonychus* Voss 1954: 211
P. titschacki Voss 1954: 211. Peru.
- Phacepholis* Horn 1876: 95
P. viridis (Champion 1911: 336) [= *P. pallidus* Pierce 1909: 361, not available; *P. pallidulus* (Emden 1936: 28); *P. eximus* (Buchanan 1939: 27)]. *Mexico.
- Platymus* Sahlberg 1823: 29 [= *Platymus* Schoenherr 1823: col. 1140, not Sahlberg 1823;
Pseudocyphus Schaeffer 1905: 179; *Eustylomorphus* Pierce 1915: 9; *Pachyus* Kuschel 1955: 305]
P. agonista (Germar 1824: 433). Brazil.
P. angustifrons (Voss 1934: 68). Costa Rica.
P. atrosignatus Lucas 1857: 155 [= *P. sigillatus* (Pascoe 1880: 421)]. Argentina, Brazil, and Peru.
P. besckei (Germar 1824: 436) [= *P. besckii* Germar 1824: 436, incorrect original spelling].
Brazil.

- P. bruchi* Hustache 1926: 166. Argentina.
- P. cavirostris* Boheman 1840: 167. Brazil.
- P. crassicornis* Lucas 1857: 156. Peru.
- P. dianae* Boheman 1840: 169 [= *P. danae* Jacob 1937: 157, lapsus]. Argentina, Brazil, and Paraguay.
- P. duponti* Boheman 1833: 633. Brazil.
- P. effusus* (Pascoe 1880: 421). Brazil, Ecuador, and Peru.
- P. elegantulus* Hustache 1923: 290. Argentina, Bolivia, Paraguay, and Uruguay.
- P. eustalooides* (Champion 1911: 284) [= *P. subvittatus* (Voss 1940: 14), not Voss 1932], **syn. nov.**; *P. vossi* Kuschel, 1955: 309, **syn. nov.**. Brazil.
- P. fasciatus* Boheman 1840: 166. Brazil.
- P. gratiosus* (Champion 1911: 285). Brazil.
- P. gyllenhali* Boheman 1840: 158. South America.
- P. humilis* (Erichson 1847: 129) [= *P. squamipunctatus* (Pierce 1915: 9)]. Peru.
- P. hystricosus* (Germar 1824: 434). Brazil.
- P. latacungae* Kirsch 1889: 17. Brazil and Ecuador.
- P. marmoratus* (Marshall 1922: 185). Brazil and Guyana.
- P. mollis* Faust 1890: 175. Argentina and Paraguay.
- P. nodipennis* Sahlberg 1823: 29 [= *P. cultricollis* Germar 1824: 432; *P. faldermanni* Rosenschoeld 1840: 157]. Argentina, Brazil, and Paraguay.
- P. ochraceus* (Hustache 1938: 280). Bolivia and Brazil.
- P. perlepidus* Boheman 1833: 636. Argentina, Brazil, and Paraguay.
- P. piscatorius* (Germar 1824: 434). Brazil.
- P. planicollis* Hustache 1926: 165. Argentina.
- P. prasinus* Boheman 1833: 634. Argentina, Brazil, and Paraguay.
- P. sellatus* (Marshall 1922: 187). Brazil.
- P. septempunctatus* Boheman 1833: 637 [= *P. novemmaculatus* Boheman 1833: 638; *P. septemmaculatus* Boheman 1833: 638; *P. mutabilis* Boheman 1840: 171]. Brazil.
- P. signatus* Rosenschoeld 1840: 172. Brazil.
- P. silbermanni* Rosenschoeld 1840: 162. Brazil and Venezuela.
- P. transversesignatus* Boheman 1840: 165. Brazil.
- P. tripunctatus* Kirsch 1874: 389. Bolivia and Peru.
- P. undulatus* Boheman 1833: 635. Brazil.
- P. vermiculatus* (Hustache 1938: 279). Brazil.
- P. viridis* (Voss 1934: 67). Brazil.
- P. wahlenbergii* Boheman 1840: 159. Guyana.
- Plectrophoroides* Wibmer & O'Brien 1986: 50 [= *Plectrophorus* Schoenherr 1826: 106, not Féruccac 1819]
- P. acuminatus* (Chevrolat 1879: LXXVII). Colombia.
- P. albilabris* (Chevrolat 1879: LXXVIII). Colombia.
- P. bifasciatus* (Chevrolat 1879: LXXVII). Bolivia and Colombia.
- P. humeralis* (Chevrolat 1879: CXLVIII). Argentina and Brazil.
- P. impressicollis* (Chevrolat 1879: LXXVII). Brazil, French Guiana, and Guyana.
- P. lutra* (Schoenherr 1826: 106) [= *P. serrimanus* (Gyllenhal 1840: 109)]. Brazil, French Guiana, Suriname, and Venezuela.
- P. quadrimaculatus* (Chevrolat 1881: XXXVIII). Bolivia, Brazil, and Peru.

Priocyphopsis Lanteri 1990: 415

P. humeridens (Hustache 1926: 162). Argentina.

Priocyphus Hustache 1939 [= *Prionocyphus* Blackwelder 1947: 795, lapsus]

P. bosqi Hustache 1939: 43. Argentina.

P. hustachei (Kuschel 1950: 548). Argentina.

P. inops (Kuschel 1950: 549). Argentina.

P. kuscheli Lanteri 1990: 412. Paraguay.

Pycnophilus Sharp 1891: 166 (**new placement**, R. Anderson, personal comm.)

P. piceus Sharp 1891: 167. Costa Rica.

Rhynchuchus Boheman 1859: 125

R. acuminatus Boheman 1859: 125. Ecuador.

Stenocyphus Marshall 1922: 183

S. bituberosus (Gyllenhal 1833: 622). Brazil.

Symmathetes Schoenherr 1847: 31

S. kollaris Schoenherr 1847: 31. Brazil.

S. setulosus Hustache 1923: 286. Argentina, Bolivia, Brazil, and Paraguay.

Teratopactus Heller 1921: 20

T. capucinus (Perty 1832: 73) [= *T. senex* (Boheman 1833: 582)]. Brazil.

T. gibbicollis (Boheman 1833: 582). Brazil.

T. nodicollis (Boheman 1833: 585) [= *T. serripes* (Boheman 1840: 28); *T. angulicollis* (Lucas 1857: 153); *T. paulanus* (Fonseca & Autuori 1933: 140)]. Argentina, Brazil, and Paraguay.

T. perpastus (Boheman 1840: 29) [= *T. tuberculatus* (Hustache 1947: 63)]. Argentina, Bolivia, Brazil, Paraguay, and Uruguay.

T. retusus (Boheman 1833: 583). Brazil.

T. tuberculatus (Arrow 1903: 257). Paraguay.

Tetragonomus Champion 1911: 240

T. tuberosus Champion 1911: 240. Colombia and Panama.

Thoracocypus Emden 1936: 81

T. denticollis Emden 1936: 81. Brazil.

T. hatschbachi Bondar 1949: 174. Brazil.

Trichaptus Pascoe 1880: 422

T. mutillarius (Perty 1832: 70) [= *T. myrmosarius* (Perty 1832: 71); *T. linnei* (Boheman 1833: 629); *T. linnaei* Schoenherr 1840: 153, lapsus]. Brazil.

Trichocyphus Heller 1921: 21 [= *Tricocyphus* Hustache 1939: 43, lapsus; *Squamamitrus* Voss 1947: 48]

T. formosus (Erichson 1834: 258) [= *T. rubricollis* (Blanchard 1847: 202); *T. nitidicollis* (Voss 1947: 46); *T. pulcher* Kuschel 1949: 11]. Argentina, Bolivia, Chile, and Peru.

Trichonaupactus Hustache 1939: 45

T. densius Hustache 1939: 46 [= *T. albescens* Hustache 1939: 47]. Argentina.
T. rexus Hustache 1939: 47. Uruguay.

Saurops Kuschel 1955: 306
S. cavifrons (Hustache 1926: 169). Argentina.

Wagneriella Hustache 1923: 294
W. lineata Hustache 1923: 295 [= *W. albula* Hustache 1923: 296]. Argentina, Bolivia, and Uruguay.

Acknowledgments. I would like to thank Bob Anderson, Anne Howden, Charlie O'Brien, and Guillermo Wibmer for their useful comments on the manuscript.

Literature cited

- BORDÓN, C. 1991. El género *Macrostylus* Boheman (Col. Curc. Brachyderinae, Naupactini) en Venezuela. *Acta Biologica Venezolana* 13(1-2): 1-50.
- BORDÓN, C. 1997. El género *Naupactus* Dejean (Coleoptera: Curculionidae) en Venezuela. *Acta Biologica Venezolana* 17(2): 11-51.
- BRIGHT, D. E. 1994. Revision of the genus *Sitona* (Coleoptera: Curculionidae) of North America. *Annals of the Entomological Society of America* 87(3): 277-306.
- CRACRAFT, J. 1992. The species of the birds-of-paradise (Paradisaeidae): Applying the phylogenetic species concept to a complex pattern of diversification. *Cladistics* 8: 1-43.
- HOWDEN, A. T. 1986. The *Pandeleteius armatus* group, with description of a new species from Mexico (Coleoptera: Curculionidae). *Coleopterists' Bulletin* 40(4): 317-323.
- HOWDEN, A. T. 1993a. *A catalog of the Coleoptera of America North of Mexico. Family: Curculionidae. Subfamily: Polydrosinae. Tribe: Tanymecini.* Agriculture Handbook 529-143e, United States Department of Agriculture, Washington D.C.
- HOWDEN, A. T. 1993b. Revision of the Mesoamerican species of *Airosimus* Howden (Coleoptera: Curculionidae). *Coleopterists' Bulletin* 47(3): 249-267.
- HOWDEN, A. T. 1996. Neotropical *Pandeleteius* (Coleoptera: Curculionidae) with irregular striae. *Canadian Entomologist* 128: 877-955.
- HOWDEN, A. T. 1998. Review of the *Pandeleteius biseriatus* species-group (Coleoptera: Curculionidae), with description of a new species. *Canadian Entomologist* 130: 367-375.
- KUSCHEL, G. 1995. A phylogenetic classification of Curculionoidea to families and subfamilies. *Memoirs of the Entomological Society of Washington* 14: 5-33.
- LANTERI, A. A. 1990a. Revisión sistemática y análisis filogenético de las especies del género *Enoplopactus* Heller 1921 (Coleoptera: Curculionidae). *Boletín de la Sociedad de Biología de Concepción (Chile)* 61: 71-92.
- LANTERI, A. A. 1990b. Systematic revision and cladistic analysis of *Phacepholis* Horn (Coleoptera: Curculionidae). *Southwestern Entomologist* 15(2): 179-204.
- LANTERI, A. A. 1990c. Revisión sistemática del género *Cyrtomon* Schoenherr (Coleoptera: Curculionidae). *Revista Brasileira de Entomologia* 34(2):387-402.

- LANTERI, A. A. 1990d. Revisión sistemática del género *Priocyphus* Hustache 1939 y creación de los géneros *Priocyphopsis* y *Lamprocyphopsis* (Coleoptera, Curculionidae). *Revista Brasileira de Entomologia* 34(2): 403-422.
- LANTERI, A. A. 1992. Systematics, cladistics and biogeography of a new weevil genus, *Galapaganus* (Coleoptera: Curculionidae) from the Galapagos islands, and coasts of Ecuador and Peru. *Transactions of the Entomological Society of America* 118(2): 227-267.
- LANTERI, A. A. 1995. Systematic revision of *Erycideus* Pascoe (Coleoptera: Curculionidae). *Entomologica Scandinavica* 26: 393-424.
- LANTERI, A. A. & N. B. DÍAZ. 1994. Systematic study and cladistic analysis of the genus *Aramigus* Horn (Coleoptera: Curculionidae). *Transactions of the American Entomological Society* 120(2): 113-144.
- LANTERI, A. A. & J. J. MORRONE. 1991. Cladistic analysis of *Priocyphus* Hustache and related genera (Coleoptera: Curculionidae). *Proceedings of the Entomological Society of Washington* 93(2): 278-287.
- LANTERI, A. A. & J. J. MORRONE. 1995. Cladistic analysis of the *Naupactus leucoloma* species group, *Eurymetopus*, and *Atrichonotus* (Coleoptera: Curculionidae). *Revista de la Sociedad Entomológica Argentina* 54(1-4): 99-112.
- LANTERI, A. A. & C. W. O'BRIEN. 1990. Taxonomic revision and cladistic analysis of *Atrichonotus* Buchanan (Coleoptera: Curculionidae). *Transactions of the American Entomological Society* 116(3): 697-725.
- MARVALDI, A. E. 1997. Higher level phylogeny of Curculionidae (Coleoptera: Curculionoidea) based mainly on larval characters, with special reference to broad-nosed weevils. *Cladistics* 13: 285-312.
- MARVALDI, A. E. 1998. Larvae of Entiminae (Coleoptera: Curculionidae): Tribal diagnoses and phylogenetic key, with a proposal about natural groups within Entimini. *Entomologica Scandinavica* 29: 89-98.
- MAY, B. M. 1993. Larvae of Curculionoidea (Insecta: Coleoptera): A systematic overview. *Fauna of New Zealand* 28: 1-221.
- MAY, B. M. 1994. An introduction to the immature stages of Australian Curculionoidea. In: E.C. Zimmerman. *Australian weevils, volume II, Brentidae, Eurhynchidae, Apionidae and a chapter on immature stages by Brenda May*. Centre of Scientific and Industrial Research, Melbourne.
- MORRIS, M. G. 1995. Recent advances in the higher systematics of Curculionoidea, as they affect the British fauna. *Coleopterist* 4(1):21-30.
- MORRONE, J. J. 1996. Clave ilustrada para la identificación de las familias sudamericanas de gorgojos (Coleoptera: Curculionoidea). *Revista Chilena de Entomología* 23: 59-63.
- MORRONE, J. J. 1997. Argentinean weevils (Coleoptera: Curculionidae): Preliminary overview, with nomenclatural and distributional notes. *Physis* (Buenos Aires), C 53(124-125): 1-17.
- MORRONE, J. J. 1998. *Anomophthalmina subtrib. n.*, an endemic subtribe of Entimini (Coleoptera: Curculionidae) from western Patagonia. *Revista de la Sociedad Entomológica Argentina* 57(1-4): 85-89.
- MORRONE, J. J. & S. ROIG-JUÑENT. 1995. *The diversity of Patagonian weevils: An illustrated checklist of the Patagonian Curculionoidea (Insecta: Coleoptera)*. Literature of Latin America, Buenos Aires.

- NORMARK, B. B. & A. A. LANTERI. 1996. *Aramigus uruguayensis* (Coleoptera : Curculionidae), a new species based on mitochondrial DNA and morphological characters. *Entomological News* 107(5) : 311-316.
- O'BRIEN, C. W. & G. J. WIBMER. 1982. Annotated checklist of the weevils (Curculionidae *sensu lato*) of North America, Central America, and the West Indies (Coleoptera: Curculionoidea). *Memoirs of the American Entomological Institute* 34: 1-382.
- O'BRIEN, C. W. & G. J. WIBMER. 1984. Annotated checklist of the weevils (Curculionidae *sensu lato*) of North America, Central America, and the West Indies (Coleoptera: Curculionoidea). Supplement 1. *Southwestern Entomologist* 9(3): 286-307.
- TANNER, V. M. 1987. Notes on American *Sitona* (Coleoptera: Curculionidae), with three new species. *Great Basin Naturalist* 47(1): 168-174.
- THOMPSON, R. T. 1992. Observations on the morphology and classification of weevils (Coleoptera, Curculionoidea) with a key to major groups. *Journal of Natural History* 26: 835-891.
- WIBMER G. J. & C. W. O'BRIEN. 1986. Annotated checklist of the weevils (Curculionidae *sensu lato*) of South America (Coleoptera: Curculionoidea). *Memoirs of the American Entomological Institute* 39: 1-563.
- WIBMER G. J. & C. W. O'BRIEN. 1989. Additions and corrections to annotated checklists of the weevils of North America, Central America, and the West Indies, and of South America. *Southwestern Entomologist* 13: 1-49.
- ZIMMERMAN, E. C. 1994. *Australian weevils. Volume I. Anthribidae to Attelabidae*. Centre of Scientific and Industrial Research, Melbourne.